

RESEARCH

A comparison of complications between Laparoscopic appendicectomy and Open appendicectomy.

DOI: 10.15436/JAS.2.1.2

Shashirekha CA¹ ; Ravikiran HR² ; Sankar Vikas³ ; Prasad Krishna⁴ ; Sreeramulu PN⁵ ; Sathanantham Dinesh Kumar⁶

1. Asso. Prof., Dept. of General Surgery

2. Asst. Prof., Dept. of General Surgery

3. Post Graduate, Dept. of General Surgery

4. Prof., Dept. of General Surgery

5. Prof., Dept. of General Surgery

6. Post Graduate., Dept. of General Surgery

Dept. of General Surgery, R.L.Jalappa Hospital & Research Institute, Attached to Sri Devaraj Urs Medical College, Tamaka, Kolar.

RECEIVED DATE: 25-11-2016; ACCEPTED DATE: 06-01-2016; PUBLISHED DATE: 20-01-2017

CORRESPONDENCE AUTHORS: Sathanantham Dinesh Kumar

E-MAIL: dinesh.sathanantham@gmail.com

CONFLICTS OF INTEREST

THERE ARE NO CONFLICTS OF INTEREST FOR ANY OF THE AUTHORS.

ABSTRACT:

Laparoscopic appendicectomy was first described by Semm in 1982. Since the time of its inception it has struggled to prove its supremacy over open technique. This is due to the fact that open appendicectomy is done by a small right iliac fossa incision most of the times and postoperative period is usually uneventful. While the complications that occur with both the techniques are same i.e. superficial or deep surgical site infections, the incidence of these infections is hypothesized to be less with laparoscopic appendicectomy. There are a number of randomized studies, meta-analyses and systematic reviews comparing the two techniques but the heterogeneity of the measured variables and other weaknesses in the methodology have not allowed to draw definitive conclusions and generalizations.

INTRODUCTION

Laparoscopic appendicectomy was first described by Semm in 1982. Since the time of its inception it has struggled to prove its supremacy over open technique. This is due to the fact that open appendicectomy is done by a small right iliac fossa incision most of the times and postoperative period is usually uneventful. While the complications that occur with both the techniques are same i.e. superficial or deep surgical site infections, the incidence of these infections is hypothesized to be less with laparoscopic appendicectomy. There are a number of randomized studies, meta-analyses and systematic reviews comparing the two techniques but the heterogeneity of the measured variables and other weaknesses in the methodology have not allowed to draw definitive conclusions and generalizations.

OBJECTIVES

To compare the incidence of complications with laparoscopic and open appendicectomies.

MATERIALS AND METHODS

A retrospective analysis of patient data compiled by assessment of follow up recorded over the period encompassing from January 2010 – July 2016 at RL Jalappa hospital and research centre, in department of General surgery, Tamaka, Kolar. The comparison of the incidence of complications with the two procedures was done using logistic regression model.

Inclusion criteria

Subjects who underwent appendicectomy during the study period in our center.

Exclusion criteria


Subjects operated on emergency basis for appendicular perforation or abscess.

Subjects with stump appendicitis

RESULTS

A total of 469 subjects underwent appendicectomy during the study period of which 103 subjects underwent laparoscopic surgery while 366 subjects underwent open procedure. Out of the 103 subjects who underwent laparoscopic surgery, 49 were male and 54 were female and out of the 366 subjects who underwent open surgery, 222 were male and 144 were female.

Figure 1



One of the open appendicectomies mentioned above was started as laparoscopic procedure and had to be converted to open procedure due to mass formation.

Out of the 103 subjects who underwent laparoscopic procedure, 54 had recurrent appendicitis and 43 had acute appendicitis. While one subject had appendicular perforation, 5 subjects underwent diagnostic laparoscopy for chronic pain in abdomen and on identification of appendicular pathology, appendicectomy was done in the same sitting.

Out of the 366 subjects who underwent open procedure, 77 had recurrent appendicitis and 243 had acute appendicitis. 35 subjects had appendicular perforation. 7 subjects underwent interval appendicectomy for resolved appendicular mass. In one subject diagnostic laparoscopy for chronic pain in abdomen was taken up and subsequently appendicular mass was detected and had to be converted into open procedure as the mass could not be dissected laparoscopically.

Table 1. Indications for the procedures

Indication	Laparoscopic procedure(n=103)	Open procedure(n=366)
Recurrent appendicitis	54 (52.4%)	77(21.0%)
Acute appendicitis	43(11.7%)	243(66.3%)
Appendicular perforation	1(0.009%)	35(0.09%)
Chronic pain in abdomen	5(0.04%)	1*(0.0027%)

*started as diagnostic laparoscopy and later open appendicectomy was done.

A total of 74 subjects developed complications in the open appendicectomy group which accounts to 20.2% of the 366 subjects while a total of 20 complications were noted in the laparoscopic appendicectomy group which accounts to 19.4% of the 103 subjects. No deaths were observed in either of the groups.

Table 2. Overall complication rates

Complications	Open appendicectomy (n=366)	Laparoscopic appendicectomy (n=103)
Total	74 (20.2%)	20 (19.4%)

All the subjects who developed complications were treated by regular dressings and antibiotics according to culture sensitivity of discharge followed by secondary closure in case of superficial surgical site infections and ultrasonogram guided aspiration and antibiotics according to culture sensitivity of aspirate in case of deep surgical site infections.

DISCUSSION

Acute appendicitis is the most common general surgical emergency encountered and early surgical intervention improves the outcomes. Richard Hall reported the first survival of a patient after removal of perforated appendix and later in 1894 McBurney described the right lower quadrant muscle splitting incision for appendicectomy. It was almost a century before laparoscopic appendicectomy was described in 1982. Laparoscopic appendicectomy is being offered in our center for almost a decade now but it is limited by the cost factor. Also it does not offer any major advantages over open procedure as the scar is small and length of hospital stay is usually the same with both the techniques. In this scenario the study of incidence of complications with these two is warranted.

In this retrospective study we found that the incidence of complications in subjects who underwent open appendicectomy was 20% and the incidence of complications in subjects who underwent laparoscopic appendicectomy as 18%. In 5% of the subjects procedure started as laparoscopic technique but later converted to open due to various reasons.

Thus we say that laparoscopic appendicectomy has minimal or no advantage over open procedure and is not economical to the patient, based on this study conducted in this teaching hospital.

However this study has various limitations like smaller sample size of laparoscopic group due to monetary issues and less awareness among people, ours being a rural setup and the study itself being a single centric, the results cannot be generalised to the entire population. Furthermore the rate of complications and incidence of conversion of laparoscopic to open procedure differs from surgeon to surgeon.

CONCLUSIONS

Laparoscopic appendicectomy has slightly less incidence of complications while being more costly and thus offers little benefit as compared to open appendicectomy. However further studies with larger sample sizes and multicentric trials within and across the country are needed to generalise these results to the world population.

REFERENCES

1. Katkhouda Namir, Mason Rodney J, Towfigh Shirin, Gevorgyan Anna, Essani Rahila. Laparoscopic versus Open appendectomy- A prospective randomized double-blind study. *Ann Surg* 2005; 242:439-450.
2. Semm K. Endoscopic appendectomy. *Endoscopy*. 1983;15:59-64.
3. Majeed AW, Troy G, Nicholl JP, et al. Randomized, prospective, singleblind comparison of laparoscopic versus small incision cholecystectomy. *Lancet*. 1996;347:989-994.
4. McBurney C. The incision made in the abdominal wall in case of appendicitis with a description of a new method of operating. *Ann Surg*. 1894;20-38.
5. Guller U, Hervey S, Purves H, et al. Laparoscopic versus open appendectomy: outcomes comparison based on a large administrative database. *Ann Surg*. 2004;239:43-52.
6. Attwood SE, Hill AD, Murphy PG, et al. A prospective randomized trial of laparoscopic versus open appendectomy *Surgery*. 1992;112:497-501.
7. Cox MR, McCall JL, Tooouli J, et al. Prospective randomized comparison of open svuesr laparoscopic appendectomy in men. *World J Surg*. 1996;20:263-266.
8. Frazee RC, Roberts JW, Symmonds RE, et al. A prospective randomized trial comparing open versus laparoscopic appendectomy. *Ann Surg*. 1994;219:725-728.
9. Hansen JB, Smithers BM, Schache D, et al. Laparoscopic versus open appendectomy: prospective randomized trial. *World J Surg*. 1996;20:17-20.
10. Heikkinen TJ, Haukipuro K, Hulkko A. Cost effective appendectomy: open or laparoscopic? a prospective randomized study. *Surg Endosc*. 1998;12:1204-1208.
11. Hellberg A, Rudberg C, Kullman E, et al. Prospective randomized multicentre study of laparoscopic versus open appendectomy. *Br J Surg*. 1999;86:48-53.
12. Ignacio RC, Burke R, Spencer D, et al. Laparoscopic versus open appendectomy: what is the real difference? Results of a prospective randomized doubleblinded trial. *Surg Endosc*. 2004;18:334-337.
13. Kazemier G, De Zeeuw GR, Lange J, Fet al. Laparoscopic vs. open appendectomy: a randomized clinical trial. *Surg Endosc*. 1997;11:336-340.
14. Klingler A, Henle K,P Beller S, et al. Laparoscopic appendectomy does not change the incidence of postoperative infectious complications. *Am J Surg*. 1998;175:232-235.
15. Kum CK, Ngoi SS, Goh PM, et al. Randomized controlled trial comparing laparoscopic and open appendectomy. *Br J Surg*. 1993;80:1599-1600.
16. Laine S, Rantala A, Gullichsen R, et al. Laparoscopic appendectomy: is it worthwhile? a prospective, randomized study in young women. *Surg Endosc*. 1997;11:95-97.
17. Larsson PG, Henriksson G, Olsson M, et al. Laparoscopy reduces unnecessary appendicectomies and improves diagnosis in fertile women: a randomized study. *Surg Endosc*. 2001;15:200-202.
18. Long KH, Bannon MP, Zietlow SP, et al. Laparoscopic Appendectomy Interest Gropu: a prospective randomized comparison of laparoscopic appendectomy with open appendectomy: clinical and economic analyses. *Surgery*. 2001;129:390-400.
19. Macarulla E, Vallet J, Abad JM, et al. Laparoscopic versus open appendectomy: a prospective randomized trial. *Surg Laparosc Endosc*. 1997;7:335-339.
20. Martin LC, Puente I, Sosa JL, et al. Open versus laparoscopic appendectomy: a prospective randomized comparison. *Ann Surg*. 1995;222:256-261.
21. Milewczyk M, Michalik M, Ciesielski M. A prospective, randomized, unicenter study comparing laparoscopic and open treatments of acute appendicitis. *Surg Endosc*. 2003;17:1023-1028.
22. Minne L, Varner D, Burnell A, et al. Laparoscopic vs. open appendectomy: prospective randomized study of outcomes. *Arch Surg*. 1997;132:708-711.
23. Mutter D, Vix M, Bui A, et al. Laparoscopy not recommended for routine appendectomy in men: results of a prospective randomized study. *Surgery*. 1996;120:71-74.
24. Ortega AE, Hunter JG, Peters JH, et al. A prospective, randomized comparison of laparoscopic appendectomy with open appendectomy. *Am J Surg*. 1995;169:208-212.
25. Pedersen AG, Petersen OB, Wara P, et al. Randomized clinical trial of laparoscopic versus open appendectomy *Br J Surg*. 2001;88:200-205.

26. Reiertsen O, Larsen S, Trondsen E, et al. Randomized controlled trial with sequential design of laparoscopic versus conventional appendectomy. *Br J Surg*. 1997;84:842–847.
27. Tate JJ, Dawson JW, Chung SC, et al. Laparoscopic versus open appendectomy: prospective randomised trial. *Lancet*. 1993;342:633–637.
28. Chung RS, Rowland DY, Li P, et al. A metaanalysis of randomized controlled trials of laparoscopic versus conventional appendectomy. *Am J Surg*. 1999;177:250–256.
29. Garbutt JM, Soper NJ, Shannon WD, et al. Metaanalysis of randomized controlled trials comparing laparoscopic and open appendectomy. *Surg Laparosc Endosc*. 1999;9:17–26.
30. Golub R, Siddiqui F, Pohl D. Laparoscopic versus open appendectomy: a metaanalysis. *Am Coll Surg*. 1998;186:545–553.
31. Sauerland S, Lefering R, Holthausen U, et al. Laparoscopic vs conventional appendectomy: a metaanalysis of randomised controlled trials. *Arch Surg*. 1998;383:289–295.
32. Fingerhut A, Millat B, Borrie F. Laparoscopic versus open appendectomy: time to decide. *World J Surg*. 1999;23:835–845.
33. McCall JL, Sharples K, Jadallah F. Systematic review of randomized controlled trials comparing laparoscopic with open appendectomy. *Br J Surg*. 1997;84:1045–1050.
34. Sauerland S, Lefering R, Neugebauer EA. Laparoscopic versus open surgery for suspected appendicitis. *Cochrane Database Syst Rev*. 2002;(1):CD001546.
35. Slim K, Pezet D, Chipponi J. Laparoscopic or open appendectomy? critical review of randomized, controlled trials. *Dis Colon Rectum*. 1998;41:398–403.
36. Heinzelmann M, Simmen H, CPummins AS, et al. Is laparoscopic appendectomy the new “gold standard”? *Arch Surg*. 1995;130:782–785.
37. Tate JJ, Chung SC, Dawson J, et al. Conventional versus laparoscopic surgery for acute appendicitis. *Br J Surg*. 1993;80:761–764.
38. Vallina VL, Velasco JM, McCulloch CS. Laparoscopic versus conventional appendectomy. *Ann Surg*. 1993;218:685–692.
39. LeLorier J, Gregoire G, Benhaddad A, et al. Discrepancies between metaanalyses and subsequent large randomized, controlled trials. *N Engl J Med*. 1999;337:536–544.
40. Katkhouda N, Friedlander MH, Grant SW, et al. Intraabdominal abscess rate after laparoscopic appendectomy. *Am J Surg*. 2000;180:456–459.
41. Paik PS, Towson JA, Anthone GJ, et al. Intraabdominal abscesses following laparoscopic and open appendectomies. *J Gastrointest Surg*. 1997;1:188–193.
42. Lord RV, Sloane DR. Early discharge after open appendectomy. *Aust N Z J Surg*. 1996;66:361–365.
43. Rao PM, Rhea JT, Novelline RA, et al. Effect of computed tomography of the appendix on treatment of patients and use of hospital resources. *N Engl J Med*. 1998;338:141–146.
44. Enochsson L, Hellbegr A, Rudberg C, et al. Laparoscopic vs. open appendectomy in overweight patients. *Surg Endosc*. 2001;15:387–392.