

Osmotic Dehydration and Assessment of Quality Attributes of Seasonal Vegetable Crops: Carrot and Beetroot Cubes

DOI: 10.25177/JFST.5.4.RA.10658

Research

Accepted Date: 04th August 2020; Published Date: 10th August 2020

Copy rights: © 2020 The Author(s). Published by Sift Desk Journals Group

This is an Open Access article distributed under the terms of the Creative Commons Attribution License

[\(http://creativecommons.org/licenses/by/4.0/\)](http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.**Shamsher Ali^{1*}, Alka Sharma¹**

¹ Department of Food Technology, Guru Jambheshwar University of Science and Technology, Hisar, Haryana-125001, India

CORRESPONDENCE AUTHOR

Shamsher Ali

Email: shamsher.gjust@gmail.com

CITATION

SHAMSHER ALI, Osmotic Dehydration and Assessment of Quality Attributes of Seasonal Vegetable Crops: Carrot and Beetroot Cubes (2020) Journal of Food Science & Technology 5(4) pp:199-217

ABSTRACT

Fulfilling the food demand of a growing population is the biggest challenge as lot of food globally got wasted due to improper storage and processing. Osmotic dehydration offers high-quality preservation and maintenance of the integrity of fruits & vegetables. Response surface methodology was performed to estimate the main effect of osmotic dehydration process on quality attributes of carrot and beetroot cubes. Higher values of the osmotic solution of salt and sucrose at sample to solution ratio of 1:5 had provoked higher flows of water and solutes through the carrot and beetroot cubes. The range of NaCl concentration varies from 4-12 % w/v in carrot and 12-16 % w/v in beetroot for 2, 4 and 6 hour. However, sucrose concentration varies from 40-60° Brix in both carrot and beetroot. Quality attribute of carrot and beetroot including ascorbic acid, carotenoid, total phenol etc. didn't changed on recommended process variables 50° Brix of sucrose + 8 % w/v sodium chloride for carrot and 50° Brix of sucrose + 14 % w/v NaCl for beetroot under osmotic dehydration for 4 hours. It was considered to get maximum water loss, weight reduction, subsequent rehydration ratio, overall acceptability and minimum solute gain of rehydrated product.

Keywords: Osmotic dehydration; Carrot; Beetroot; Preservation; Response Surface Methodology

1. INTRODUCTION

India has been bestowed with a broad range of climate and physio-geographical conditions and is best suitable for horticulture crops like fruits, vegetables, nuts, flowers, and plantation crops (Gupta et al., 2019). Fruits and vegetables contribute an important source of nutrients in the daily human diet in accordance, to the horticulture database 2018, the globe fruits production are estimated to 184 million metric tons (MMT) and vegetables 97 MMT. India is that the second-largest producer of fruits and vegetables and its annual production is 89 and 163 MMT respectively during 2017-2018 contributing 11.2% of world production (Handbook on Horticulture Statistic, 2018). Despite the high contribution of India to global food production, approx. 18% of the whole fruits and vegetables got wasted thanks to improper storage and processing (Balaji and Arshinder, 2016).

Carrot and beetroot are the most significant seasonal root vegetables grown extensively in the tropical region during winter seasons. Carrots (*Daucus carota*) have significant multidimensional applications including preparation of salad, cooked vegetables, stews, curries, sweetmeat, juices, fermented pickles, flakes, and soups. Carrots are rich in nutrients like carotenoids, vitamins (B1, B2, and B6), and minerals (Singh et al., 2012). The antioxidant properties of beta carotene in carrots help to weigh down the aging process (Singh et al., 1996), reduce cholesterol production, and thus minimize the danger of heart diseases (Di Pietro et al., 2016). Falcarinol, a compound presents in carrots is additionally reported to stop breast, lung, and colon cancers (Tan et al., 2014; Kobæk-Larsen et al., 2018). Vitamins and minerals of carrots detoxify the liver and reduce decay and damage (Pryor et al., 2000). vegetable commonly referred to as beetroot may be a species of phanerogam of the goosefoot family. Beetroot is peeled off, steamed, eaten warm with butter, cooked, pickled, and eaten cold as a condiment, shredded raw or eaten as a salad. Beetroots also are high in folate, solvable and insoluble dietary fiber and antioxidants. Betalains in the juice of beetroot act as antioxidants to stop oxidative stress and also help to relax the mind (Clifford et al., 2017). The betacyanin present in beetroot additional-

ly to its detoxification activity is additionally reported to own chemotherapeutic activity in numerous kinds of cancers (Kapadia et al., 2011). Dietary consumption of beetroot nitrate increases the extent of alpha-lipoprotein and consequently reduces the beta-lipoprotein and minimizes the danger of coronary diseases (Singh et al., 2015). It increases somatic cell count so it's an awfully good option for anemic people because it is extremely rich in iron content (Hatlin Sugi, 2014). Moreover, beetroots purify the blood and strengthen the system against common diseases including jaundice, hepatitis, nausea, and vomiting.

Preservation of fruits, vegetables, and other perishable food is important for increasing its availability to satisfy the demand of the growing population; maintenance of its quality, taste, and integrity is additionally a challenging job. Numerous process technologies including freezing, canning, dehydration, etc. are used at industrial scale for the preservation of food products but it's very expensive. Therefore, there's a necessity for the straightforward and cheap alternate process which incorporates a low capital investment and offers a technique to save lots of highly perishable products and increases its availability within the region off from the assembly zones (Shi and Le Maguer, 2002). Among others, osmotic dehydration (OD) offers high-quality preservation and maintenance of the integrity of fruits & vegetables by a way of water removal without activity and reduction of microbial growth (Bahmani et al., 2016). Moreover, OD also improves nutritional value, sensory traits, and functional properties of food without changing its integrity. OD process within the food industry has several advantages like increased sugar to acid ratio, improves quality parameters, energy efficiency, packaging reduction, cost reduction, improves texture stability of pigments, better product firmness, and retention of nutrients during dehydration and storage. it's effective at ambient temperatures, that the heat damage to texture, color, and flavor is minimized.

In normal condition, fresh carrots and beetroot are stored up to three or 4 days but the period is extended up to 7 to eight months by its storage in crates cov-

ered with perforated plastic films (00 C, 93-96% relative humidity), through fermentation, pickling, canning, freeze-drying, cold storage, etc. (Madison and Coleman, 2007; Rundla and Mishra, 2018). In recent years, OD attained attention for the preservation of fruits and vegetables like banana, carrot, pineapple, mango, and leafy vegetables. It also improves to retain characteristics including color, aroma, and nutritional compositions (Akbarian et al., 2014; Mercali et al., 2011). However, the effect of osmotic dehydration on carrot and beetroot remained unaddressed. Keeping in sight the above aspects, the current study was undertaken to perform osmotic dehydration of carrot & beetroot by using different hypertonic solutions. Moreover, a comparison of varied quality attributes and drying parameters of the obtained products was done.

2. MATERIALS & METHODS


2.1 Sample Collection:-

The carrot of Hisar Garlic variety was brought from Chaudhary Charan Singh Agriculture University, Haryana, India. Beetroot obtained from the local market of Hisar, Haryana.

2.2 Osmotic dehydration of carrot and beetroot

Osmotic dehydration of carrot cubes was done in a sodium chloride solution of 4, 8, and 12 % and 40, 50, and 60° Brix of sucrose. Beetroot was osmotically dehydrated in solutions of sodium chloride: 12, 14, and 16 % and sucrose: 40, 50, and 60 ° Brix. The ratio of sample to the solution was 1:5 for 2, 4, and 6 hours to determine water loss (WL) and solute gain (SG). To obtain the maximum WL, SG, and weight reduction (WR) at optimizing temperature, fruits solution ratio, and degree Brix of the osmotic solution of sucrose was selected using Response surface methodology (RSM).

Drying is one of the oldest methods for the preservation of food by reduction of moisture content that minimize the deteriorative chemical reaction in highly perishable food. Drying of carrot and beetroot was done by two methods: Osmotic dehydration and Oven Drying.



2.3. Measurement of quality attribute and drying parameter

Determination of moisture content

Moisture content of treated as well as untreated samples was estimated just before drying by hot air oven drying method. The cubes of carrot and beetroot dried at 100°C till the constant weight of the sample was obtained and subsequently the weight of the samples were determined after cooling in desiccators. Moisture content is expressed in term of percentage (%). The loss in weight during drying was used to determine the moisture content of the sample during the OD process.

$$\text{Moisture content (\% dry basis)} = \frac{W_1 - W_2}{W_1} \times 100$$

W_1 = weight of the initial sample
 W_2 = weight of dried sample

Estimation of dry matter, water loss and solute gain.

During osmotic dehydration, WL and SG take place simultaneously. The weight of the sample is reduced due to water loss; simultaneously increase in weight due to solute permeability. Therefore, water loss is the sum of weight reduction and solute gain.

$$\text{Weight reduction (g)} = \text{WR} = (W_0 - W_t)$$

$$\text{Solute gain after OD for time t (g)} = \text{SG} = (S_t - S_0)$$

$$\text{WL} = \text{WR} + \text{SG}$$

The WL and SG during osmotic dehydration were calculated by the equation given by Ozen et al., (2002).

$$\text{WL/100g of sample} = \frac{(W_0 - W_t) + (S_t - S_0)}{W_0} \times 100$$

$$\text{SG/100g of sample} = \frac{S_t - S_0}{W_0} \times 100$$

W_0 = Initial weight of sample (g).

W_t = Weight of sample after OD at a time t (g).

S_t = Initial weight of solids (dry matter) in the sample (g).

S_0 = Weight of solids (dry matter) of the sample after OD for time t (g).

Determination of ash content

Ash content represents the inorganic residue after the destruction of organic matter. For determination of ash content, 5g of the sample was pre-weighed in clean crucible followed by heating to charring of the sample on a hot plate. The crucible with the carbon residue of ignition was placed in a muffle furnace at a temperature of 550°C until the carbon residue disappears followed by cooling and weighing. The difference in the weight by ash content was calculated by using formula

$$\text{Total Ash content (\%)} = \frac{\text{Final wt.}}{\text{Initial wt.}} \times 100$$

Rehydration ratio

Rehydration ratio (RR) was measured by soaking a 10 – 15g of each sample in sufficient volume of water (approximately 30 times the weight of sample) at room temperature (RT) (Gupta and Shukla, 2017). Rehydration was done for 12 hours, cubes reached to constant weight that were weighed after removing excess of water with the help of absorbent paper. The rehydration ratio was calculated by using formula.

$$\text{Rehydration Ratio} = \frac{W_2}{W_1}$$

W_1 = Weight of rehydrated sample

W_2 = Weight of dehydrated sample

Total phenol content

Total phenol content was determined according to Folin Ciocalteu procedure (Gonçalves et al., 2010) with slight modifications. 2g of the sample was homogenized in 80% of aqueous ethanol at a RT fol-

lowed by centrifugation at 10,000 rpm for 15 minutes, the supernatant was retained. The residue was re-extracted twice in 80% ethanol and the supernatant was evaporated to dryness by using evaporating dishes in a water bath at 40°C. The residue was dissolved in 5ml of distilled water (DW). Extract was diluted to 3 ml with distilled water followed by the addition of 0.5 ml of Folin-Ciocalteu reagent and incubation of 3 minutes. 2ml of 20% sodium carbonate was added and content was thoroughly mixed. After incubation of 60 minutes, the optical density of colored product was measured at 760 nm in UV spectrophotometer using catechol as a standard. Result was expressed as mg catechol/100g of fresh weight material. Total phenol was calculated by using formula.

$$\text{Total Phenol (mg/100g)} = \frac{M \times V \times 100}{W \times V_1 \times 100}$$

M = Concentration by graph

V = Volume conjure

V_1 = Volume of extract aliquot for color development

W = Weight of sample

Total carotenoid content

Total carotenoid content was resolute by homogenizing 1g of fresh sample in 10 ml of DW during a blender. After centrifugation, 4.5 ml of acetone was added in 0.5 ml aliquot to extract pigments. The supernatant was taken after centrifugation and absorbance was measured at 480, 645 and 663 nm by using 80% acetone as a blank within the UV spectrophotometer.

Total carotenoid was calculated by using formula.

$$\text{Carotene (g/liter)} = (\text{absorbance at 480}) - (0.114) (\text{absorbance at 663}) - (0.638) (\text{absorbance at 645})$$

Total antioxidant content using DPPH radical scavenging assay

The total antioxidant activity of the carrot and beetroot powder was estimated by using the DPPH radical scavenging protocol. DPPH solution (0.004% w/v) was prepared in 95% ethanol. a customary water-soluble vitamin solution was prepared at the concentration of 10mg/100ml specified 2 ml of freshly prepared DPPH solution (0.004% w/v) was added in

each tube. The reaction mixture was incubated within the dark for 15 min. Thereafter, the absorbance was recorded at 523 nm against the blank after 30 min. For the control, 2 ml of DPPH solution was mixed with 10 ml of ethanol. the potential of scavenging DPPH radical was calculated by using the subsequent equation (Ara and Nur, 2009).

$$\text{DPPH scavenged (\%)} = \frac{(\text{Abs.}(\text{control}) - \text{Abs.}(\text{carrot/beetroot}))}{(\text{Abs.}(\text{control}))} \times 100$$

Ascorbic acid content

The method was supported the reduction of two,6-dichlorophenol-indophenol dye by water-soluble vitamin (Shah and Nath, 2008). water-soluble vitamin content was measured on a dry weight basis.

2, 6-dichlorophenol-Indophenol Visual titration method

The dye is blue in alkaline solution and red in acid solution; reduced via water-soluble vitamin to a colorless form. The reaction is quantitative specific for water-soluble vitamin within the pH range 1-3.5.

Statistical Method

Response surface methodology (RSM) could be a very useful gizmo in product design (Box and Wilson, 1951). it's an efficient tool for the assessment of responses stricken by many factors and their interactions. RSM is reported to be an efficient tool for optimizing a process when the independent variables have a combined effect on the responses (Mudahar et al., 1989). it's an employed tool in analyzing experimental data consequent within the optimization of processes or products. it's used for designing the experiments or is also defined as an empirical statistical modeling technique employed for multivariate analysis analysis using quantitative data obtained from properly intended experiments to resolve multivariate equations simultaneously. A mathematical relation, f , was assumed for describing the correlation between each of the response variables, Y_i and also the factors x_i , such as

$$Y_i = f(x_1, x_2, x_3, \dots)$$

The exact mathematical representation of the function (f) is either unknown or extremely complex. Though, a second-order polynomial equation of the subsequent form was assumed to relate Y_{ij} and X_i .

Where, $\beta_0, \beta_i, \beta_{ii}, \beta_{ij}$ are regression coefficients. x_0, x_i, x_j are the coded independent variables linearly associated with real variables. it's accustomed fit the second or-

der polynomial equation to the experimental data.

For optimization of the OD process, the experiment was conducted per Central Composite Design (CCD) with three variables. The CCD design predict uniformly in the least constant distances from their central point (Khuri et al. Cornell J A et al.1987). the look was created by commercial statistical package, Design-Expert version 10.0.3.1 (Statease Inc., Minneapolis, USA, Trial version). The variables were salt concentration, sucrose concentration and time of osmotic dehydration process. The low to high level of NaCl concentration in carrot and beetroot varies from 4-12 % w/v and 12 to Sixteen Personality Factor Questionnaire w/v respectively. The sucrose concentration is 40-600 Brix for both carrot and beetroot. The incubation was in serious trouble 2, 4 and 6 hours. RSM used to estimate the main effect of OD process on WL, SG and WR in carrot and beetroot cube with the aim of decreasing the cost of expensive analysis methods and their associated numerical noise. (Box and Draper, 1987; Venter et al. 1996). Experiment data were randomized under Central Composite and ANOVA for response surface methodology by using the quadratic model.

3. RESULTS & DISCUSSION

Osmotic dehydration kinetics including WR, SG, WL and RR of carrot and beetroot cubes was studied using sucrose and sodium chloride salts as osmotic agents. The process parameters viz. concentration of the osmotic solution, time and sample to solution ratio were also optimized.

3.1 Proximate analysis

Proximate analysis of carrot and beetroot was carried out and results are shown in Table 3.1. The results are expressed as g/100g as on a dry weight basis. The carbohydrate, fat, protein, sucrose, and crude fibre content was reported within the range reported by Holland et al., (1991), Anon et al., (1952) and Zwart et al., (2003) respectively. The effects of practice parameters (concentration, duration of sample to solution ratio) on the kinetics of moisture loss and solute uptake during OD of carrot and beetroot cubes were accomplished according to the CCD experimental design as indicated in tables 3.4 and 3.9. The results showed that the combined effect of sugar and salt was more effective than individual concentration in case of both water loss and solute uptake during the process.

Table 3.1: Proximate analysis of carrot and beetroot

Sample	Moisture (%)	Ash (%)	Carbohydrate (%)	Fat (%)	Protein (%)	Sucrose (%)	Crude Fiber (%)
Carrot	88 ± 0.5	0.9 ± 0.2	10.4 ± 0.2	0.2 ± 0.1	0.9 ± 0.1	26 ± 1.7	2.4 ± 0.2
Beetroot	86 ± 0.5	1.1 ± 0.3	10 ± 0.2	0.3 ± 0.1	1.6 ± 0.1	78 ± 6.7	2.8 ± 0.1

Values are mean±SD of three independent determinations

3.2 Assessment of ascorbic acid, phenolic content, carotenoids and antioxidant Activity

Ascorbic acid, total carotenoid values didn't changed on any of the treatment processes. Antioxidant activity of both carrot and beetroot significantly reduced on 8% NaCl and 50° brix sucrose treatment but not on mixture of both. Total phenol level changed on single treatment of 14% NaCl and 50° brix sucrose in beet root but not carrot. On combinational treatment, no phenolic content changed in resulting carrot and beetroot (Table 3.2 and 3.3).

Table 3.2. Ascorbic Acid, Phenolic Content, Carotenoids and Antioxidant Activity of carrot

Sr. No	Sample	Fresh Sample	8% NaCl Treated Sample	50° Brix Treated Sample	8 % NaCl+ 50° Brix Sucrose treated Sample
1.	Ascorbic Acid (mg/100 g FW)	54.5 ± 0.20	50.1 ± 0.3	51.5 ± 0.4	53 ± 0.1
2.	Total Phenol (mg/100 g FW)	39.76 ± 1.3	35.2 ± 0.8	36.7 ± 0.7	39. ± 0.3
3.	Total Carotenoids (mg/1000g FW)	81 ± 1.5	77 ± 3.5	79 ± 0.5	80 ± 0.5
4.	Total Antioxidant (mg/100 g FW)	34.44 ± 0.1	27 ± 0.5***	29 ± 0.2***	32 ± 1.3

Values are mean±SD of three independent determinations

Table 3.3. Ascorbic Acid, Phenolic Content, Carotenoids and Antioxidant Activity of beetroot

Sr. No	Constituent Name	Fresh Sample	14% NaCl Treated Sample	50° Brix Scrose Treated Sample	14% NaCl+ 50° Brix Su-crose treated Sample
1.	Ascorbic Acid (mg/100 g FW)	48.4 ± 0.8	44.1 ± 0.6	46.7 ± 0.4	47.9 ± 0.5
2.	Total Phenol (mg/100 g FW)	57.64 ± 1.2	35.2 ± 0.8 ***	36.7 ± 0.7***	49. ± 0.3
3.	Total Carotenoids (mg/100 g FW)	76.81 ± 1.5	72 ± 3.5	74 ± 0.5	74 ± 0.8
4.	Total Antioxidant (mg/100 g FW)	43.12 ± 0.8	27.5 ± 0.5***	29 ± 0.2***	32 ± 1.3

Values are mean±SD of three independent determinations

Table 3.4. Observed values of dependent variables for different runs of optimization experiments for osmotic

Run	Factor 1	Factor 2	Factor 3	Response1	Response2	Response3	Response4
	A:NaCl (%)	B:Sucrose (°Brix)	C:Time (Hour)	Weight Reduction (%)	Water loss (%)	Solute gain (%)	Rehydration Ratio
1	12	40	2	25.19	30.99	5.8	2.872
2	8	50	4	36.89	50.79	12.9	2.454
3	12	60	2	28.45	39.55	11.1	2.4732
4	8	66.8179	4	35.8	49.93	13.65	2.243
5	8	50	4	37.4	49.9	12.5	2.659
6	12	60	6	35.47	48.17	12.7	2.739
7	8	50	4	37.34	49.44	12.1	2.652
8	8	50	4	37.07	48.89	11.82	2.588
9	4	40	6	22.78	29.38	6.6	2.874
10	8	50	0.63641	25.7	28.68	2.98	3.176
11	12	40	6	25.64	33.54	7.9	3.017

12	8	50	4	37.19	48.14	10.95	2.693
13	4	60	2	27.71	36.61	8.9	2.656
14	1.27283	50	4	32.7	39.39	6.68	2.833
15	14.7272	50	4	34.1	42.9	8.9	2.904
16	8	33.1821	4	21.77	27.5	5.73	3.103
17	8	50	7.36359	36.44	50.77	14.33	2.176
18	4	40	2	24.18	28.78	4.6	2.995
19	4	60	6	31.94	45.75	13.81	2.341
20	8	50	4	35.91	51.34	13.43	2.566

3.3 Mass transfer kinetics during osmotic dehydration

The detailed analyses of various responses factors for the OD of carrot and beetroot cubes are mentioned. OD process with all possible combinations, the full factorial design with three factors viz., concentration, sample to solution ratio and immersion time was used during the process of OD. The results were assessed by the multiple linear regression equation conducted for the second-order response surface model as mentioned below.

1. Mass transfer kinetics for carrot:

A.1: Fitting models for WR of carrot

Results presented in Table 3.5. The significance of each coefficient was determined through the Fischer F test and P values for the carrot. The Model F-value of 12.95 implies that the model is significant. There is only 0.02% chance that F-value will occur due to noise. Values of "Prob > F" less than 0.0500 indicate model terms are significant. In this case, B, C, A², B², C² are significant model terms. The "Lack of Fit F-value" of 30.59 implies that the Lack of Fit is significant. There is only 0.09% chance that a "Lack of Fit F-value" may occur due to noise.

Table 3.5. ANOVA of response surface quadratic model for weight reduction.

Analysis of variance table [Partial sum of squares - Type III]						
Source	Sum of Squares	df	Mean Square	F Value	p-value Prob > F	
Model	555.91	9	61.77	12.95	0.0002	significant
A-NaCl	8.06	1	8.06	1.69	0.2227	
B-Sucrose	178.51	1	178.51	37.42	0.0001	
C-Time	58.90	1	58.90	12.35	0.0056	
AB	0.020	1	0.020	4.193E-003	0.9496	
AC	2.69	1	2.69	0.56	0.4699	
BC	18.61	1	18.61	3.90	0.0765	
A ²	52.37	1	52.37	10.98	0.0078	
B ²	180.39	1	180.39	37.82	0.0001	
C ²	107.42	1	107.42	22.52	0.0008	
Residual	47.70	10	4.77			
Lack of Fit	46.19	5	9.24	30.59	0.0009	significant
Pure Error	1.51	5	0.30			
Total	603.62	19				

Final equation in terms of coded factors:

$$\begin{aligned}
 \text{WR} = & +37.56 + 0.77 * A + 3.62 * B + 2.08 * C + 0.050 * AB + 0.58 \\
 & * AC + 1.53 * BC - 2.08 * A^2 - 3.71 * B^2 - 2.90 * C^2
 \end{aligned}$$

B.1: WR response surfaces plots of carrot

Result represented in Fig. 3.1

X1=A: NaCl; X2 =B: Sucrose; Actual factor C: Time = 4




Fig. 3.1. Effect of osmotic solution of NaCl, sucrose concentration on weight reduction during osmotic dehydration of carrot cubes

A.2: Fitting models for solute gain of carrot

Results presented in Table 3.6. The significance of each coefficient was determined through the Fischer F test and P values for the carrot. The Model F-value of 10.82 implies the model is significant. There is only 0.05% chance that F-value could occur due to noise. Values of "Prob > F" less than 0.0500 indicate model terms are significant. In this case, B, C, A^2 , C^2 are significant model terms. The "Lack of Fit F-value" of 4.47 implies there is a 6.29% chance that a "Lack of Fit F-value" could occur due to noise.

The final equation in terms of coded factors:

$$\begin{aligned}
 \text{SG} = & +12.28 + 0.45 * A + 2.43 * B + 2.17 * C - 0.18 * AB - 0.40 \\
 & * AC + 0.30 * BC - 1.43 * A^2 - 0.70 * B^2 - 1.26 * C^2
 \end{aligned}$$


Table 3.6. ANOVA of response surface quadratic model for solute gain

Analysis of variance table [Partial sum of squares - Type III]						
Source	Sum of Squares	df	Mean Square	F Value	p-value	
Model	200.85	9	22.32	10.82	0.0005	significant
A-NaCl	2.71	1	2.71	1.31	0.2787	
B-Sucrose	80.37	1	80.37	38.97	< 0.0001	
C-Time	64.58	1	64.58	31.31	0.0002	
AB	0.25	1	0.25	0.12	0.7357	
AC	1.29	1	1.29	0.62	0.4477	
BC	0.73	1	0.73	0.35	0.5662	
A ²	29.65	1	29.65	14.37	0.0035	
B ²	7.15	1	7.15	3.47	0.0923	
C ²	22.85	1	22.85	11.08	0.0076	
Residual	20.63	10	2.06			
Lack of Fit	16.86	5	3.37	4.47	0.0629	significant
Pure Error	3.77	5	0.75			
Total	221.47	19				

B.2: SG response surfaces plots of carrot

Result represented in Fig. 3.2

X1=A: NaCl; X2=B: Sucrose; Actual factor C: Time = 4

**Fig. 3.2.** Effect of osmotic solution of NaCl, sucrose concentration on SG during osmotic dehydration of carrot

A.3: Fitting models for water loss of carrot

Results presented in Table 3.7. The Model F-value of 17.11 implies the model is significant. There is only a 0.01% chance that F-value could occur due to noise. Values of "Prob > F" less than 0.0500 indicate model terms are significant. In this case, B, C, A^2 , B^2 , C^2 are significant model terms.

The "Lack of Fit F-value" of 11.79 implies the Lack of Fit is significant. There is only 0.85% chance that a "Lack of Fit F-value" may occur due to noise.

Table 3.7: ANOVA for response surface quadratic model of water loss

Analysis of variance table [Partial sum of squares - Type III]						
Source	Sum of Squares	Df	Mean Square	F Value	p-value Prob > F	
Model	1390.42	9	154.49	17.11	< 0.0001	significant
A-NaCl	20.24	1	20.24	2.24	0.1653	
B-Sucrose	508.25	1	508.25	56.29	< 0.0001	
C-Time	246.84	1	246.84	27.34	0.0004	
AB	0.13	1	0.13	0.014	0.9078	
AC	0.26	1	0.26	0.028	0.8697	
BC	26.68	1	26.68	2.95	0.1164	
A^2	179.05	1	179.05	19.83	0.0012	
B^2	272.30	1	272.30	30.16	0.0003	
C^2	251.66	1	251.66	27.87	0.0004	
Residual	90.29	10	9.03			
Lack of Fit	83.23	5	16.65	11.79	0.0085	significant
Pure Error	7.06	5	1.41			
Total	1480.72	19				


The final equation in terms of coded factors:

$$\begin{aligned}
 \text{WL} = & +49.85 + 1.22 * A + 6.10 * B + 4.25 * C - 0.13 * AB + 0.18 \\
 & * AC + 1.83 * BC - 3.52 * A^2 - 4.35 * B^2 - 4.18 * C^2
 \end{aligned}$$

B.3: WL response surface plots of carrot

Result represented in Fig. 3.3

X1=A: NaCl; X2=B: Sucrose; Actual factor C: Time = 4



A.4: Fitting models for rehydration ratio of carrot

Results presented in Table 3.8. The Model F-value of 3.52 implies the model is significant. There is only 3.13% chance that F-value could occur due to noise. Values of "Prob > F" less than 0.0500 indicate model terms are significant. In this case B, C are significant model terms. The "Lack of Fit F-value" of 8.34 implies that the Lack of Fit is significant. There is only a 1.81% chance that a "Lack of Fit F-value" may occur due to noise.

Table 3.8. ANOVA of response surface quadratic model for rehydration ratio

Analysis of variance table [Partial sum of squares - Type III]						
Source	Sum of Squares	Df	Mean Square	F Value	p-value Prob > F	
Model	1.11	9	0.12	3.52	0.0313	significant
A-NaCl	9.208E-003	1	9.208E-003	0.26	0.6189	
B-Sucrose	0.66	1	0.66	18.79	0.0015	
C-Time	0.21	1	0.21	6.10	0.0331	
AB	4.763E-003	1	4.763E-003	0.14	0.7197	
AC	0.090	1	0.090	2.56	0.1404	
BC	6.698E-004	1	6.698E-004	0.019	0.8927	
A ²	0.13	1	0.13	3.63	0.0857	
B ²	8.840E-003	1	8.840E-003	0.25	0.6259	
C ²	9.614E-003	1	9.614E-003	0.28	0.6114	
Residual	0.35	10	0.035			
Lack of Fit	0.31	5	0.062	8.34	0.0181	significant
Pure Error	0.037	5	7.485E-003			
Cor Total	1.46	19				

The final equation in terms of coded factors:


$$\begin{aligned}
 \text{RR} = & +2.60 + 0.026 * A - 0.22 * B - 0.12 * C + 0.024 * AB + 0.11 \\
 & * AC - 9.150 * BC + 0.094 * A^2 + 0.025 * B^2 + 0.026 * C^2
 \end{aligned}$$

B. 4: RR response surface plots of carrot:

Result represented in Fig. 3.4

Fig. 3.4. Effect of osmotic solution of NaCl, sucrose concentration on rehydration ratio during osmotic dehydration of carrot cubes

X1=A: NaCl; X2=B: Sucrose; Actual factor C: Time = 4



2. Mass transfer kinetics for beetroot:

The effects of process parameters (concentration, duration & sample to solution ratio) on the kinetics of moisture loss and solute uptake in OD of beetroot cubes were performed according to the CCD experimental design given in Table 3.9. The CCD experimental design helps in obtaining the optimum combination of parameters for the OD of beetroot.

Table 3.9. Observed values of dependent variables for different runs of optimization experiments for osmotic dehydration of beetroot

Run	Factor 1 A:NaCl (%)	Factor 2 B:Sucrose (°Brix)	Factor 3 C:Time (Hour)	Response 1 Weight Reduction (%)	Response 2 Solute Gain (%)	Response 3 Water Loss (%)	Response 4 Rehydration Ratio
1	12	60	2	30.73	8.34	39.07	2.685
2	10.6364	50	4	33.28	9.14	42.42	2.722
3	14	50	4	34.21	10.68	44.89	2.73
4	14	50	4	35.13	9.31	44.44	2.611
5	14	50	4	33.77	11.49	45.26	2.669
6	14	50	4	34.59	11.15	45.74	2.793
7	12	60	6	27.11	11.21	38.32	2.274
8	12	40	6	25.47	6.32	31.79	2.784
9	14	33.1821	4	17.64	7.43	25.07	3.031
10	14	50	7.36359	34.89	11.17	46.06	2.091
11	16	40	2	21.33	5.99	27.32	2.771
12	16	60	6	27.54	11.56	39.1	2.643
13	16	60	2	31.77	8.61	40.38	2.32
14	14	50	4	33.76	10.44	44.2	2.733
15	17.3636	50	4	34.22	9.87	44.09	3.002
16	14	50	0.636414	19.73	4.211	23.941	3.02
17	16	40	6	25.98	6.61	32.59	2.982
18	14	66.8179	4	34.41	12.19	46.6	2.298
19	14	50	4	34.11	11.38	45.49	2.721
20	12	40	2	20.63	5.87	26.5	2.859

A.1: Fitting models for WR of beetroot

The results presented in Table 3.10. The significance of each coefficient was determined through the Fischer F test and P values for beetroot. The Model F-value of 6.57 implies the model is significant. There is only 0.35% chance that F-value will occur due to noise. Values of "Prob > F" less than 0.0500 indicate model terms are significant. In this case, B, C, B², C² are significant model terms. Values greater than 0.1000 indicate the model terms are not significant. The "Lack of Fit F-value" of 67.43 implies that the Lack of Fit is significant. There is only a 0.01% chance that a "Lack of Fit F-value" may occur due to noise.

Table 3.10. ANOVA of response surface quadratic model for weight reduction

Analysis of variance table [Partial sum of squares - Type III]					
Source	Sum of Squares	df	Mean Square	F Value	p-value
Model	559.04	9	62.12	6.57	0.0035
A-NaCl	1.33	1	1.33	0.14	0.7154
B-Sucrose	197.57	1	197.57	20.91	0.0010
C-Time	53.92	1	53.92	5.71	0.0380
AB	8.450E-003	1	8.450E-003	8.942E-004	0.9767
AC	0.080	1	0.080	8.466E-003	0.9285
BC	37.58	1	37.58	3.98	0.0741
A ²	6.77	1	6.77	0.72	0.4171
B ²	168.23	1	168.23	17.80	0.0018
C ²	126.46	1	126.46	13.38	0.0044
Residual	94.50	10	9.45		
Lack of Fit	93.12	5	18.62	67.43	0.0001
Pure Error	1.38	5	0.28		
Total	653.53	19			

The final equation in terms of coded factors:

$$\begin{aligned} \text{WR} = & +34.34 + 0.31 * A + 3.80 * B + 1.99 * C + 0.033 * AB - 0.100 \\ & * AC - 2.17 * BC - 0.69 * A^2 - 3.42 * B^2 - 2.96 * C^2 \end{aligned}$$

B. 1: WR response surface plots of beetroot

Result represented in Fig. 3.5

X1=A: NaCl; X2=B: Sucrose; Actual factor C: Time = 4




Fig. 3.5. Effect of osmotic solution of NaCl, sucrose concentration on weight reduction during osmotic dehydration of beetroot cubes

A. 2: Fitting models for SG of beetroot

The results presented in Table 3.11. The Model F-value of 9.56 implies the model is significant. There is only 0.08% chance that F-value this large will occur due to noise. Values of "Prob > F" less than 0.0500 indicate model terms are significant. In this case, B, C, B², C² are significant model terms. The "Lack of Fit F-value" of 2.36 implies that the Lack of Fit is not significant relative to the pure error. There is 18.35% chance that a "Lack of Fit F-value" this large may occur due to noise.

Table 3.11: ANOVA of response surface quadratic model for SG

Analysis of variance table [Partial sum of squares - Type III]						
Source	Sum of Squares	df	Mean Square	F Value	p-value Prob > F	
Model	94.85	9	10.54	9.56	0.0008	significant
A-NaCl	0.37	1	0.37	0.34	0.5736	
B-Sucrose	38.52	1	38.52	34.93	0.0001	
C-Time	25.31	1	25.31	22.96	0.0007	
AB	5.512E-003	1	5.512E-003	4.999E-003	0.9450	
AC	7.813E-003	1	7.813E-003	7.085E-003	0.9346	
BC	2.82	1	2.82	2.56	0.1408	
A ²	5.41	1	5.41	4.91	0.0511	
B ²	3.68	1	3.68	3.33	0.0978	
C ²	22.68	1	22.68	20.57	0.0011	
Residual	11.03	10	1.10			
Lack of Fit	7.75	5	1.55	2.36	0.1835	not significant
Pure Error	3.28	5	0.66			
Total	105.88	19				

Final equation in terms of coded factors:

$$\begin{aligned}
 \text{SG} = & +10.77 + 0.17 * A + 1.68 * B + 1.36 * C + 0.026 * AB + 0.031 * AC \\
 & + 0.59 * BC - 0.61 * A^2 - 0.51 * B^2 - 1.25 * C^2
 \end{aligned}$$

B. 2: SG response surface plots of beetroot: fig. 3.6

Result represented in Fig. 3.6

X1=A: NaCl; X2=B: Sucrose; Actual factor C: Time = 4




Fig. 3.6: Effect of osmotic solution of NaCl, sucrose concentration on SG during osmotic dehydration of beetroot cubes

A. 3: Fitting models for WL of beetroot

The results presented in Table 3.12. The Model F-value of 7.93 implies the model is significant. There is only 0.16% chance that F-value will occur due to noise. Values of "Prob > F" less than 0.0500 indicate model terms are significant. In this case, B, C, B², C² are significant model terms. The "Lack of Fit F-value" of 77.96 implies that the Lack of Fit is significant. There is only a 0.01% chance that a "Lack of Fit F-value" may occur due to noise.

Table 3.12: ANOVA of response surface quadratic model for water loss

Analysis of variance table [Partial sum of squares - Type III]						
Source	Sum of Squares	df	Mean Square	F Value	p-value Prob > F	
Model	1026.17	9	114.02	7.93	0.0016	Significant
A-NaCl	3.11	1	3.11	0.22	0.6518	
B-Sucrose	410.55	1	410.55	28.56	0.0003	
C-Time	153.12	1	153.12	10.65	0.0085	
AB	0.028	1	0.028	1.921E-003	0.9659	
AC	0.038	1	0.038	2.630E-003	0.9601	
BC	19.81	1	19.81	1.38	0.2676	
A ²	24.29	1	24.29	1.69	0.2228	
B ²	221.64	1	221.64	15.42	0.0028	
C ²	256.25	1	256.25	17.82	0.0018	
Residual	143.78	10	14.38			
Lack of Fit	141.95	5	28.39	77.96	< 0.0001	significant
Pure Error	1.82	5	0.36			
Total	1169.95	19				

The final equation in terms of coded factors:

$$\begin{aligned}
 \text{WL} = & +45.11 + 0.48 * A + 5.48 * B + 3.35 * C + 0.059 * AB - 0.069 \\
 & * AC - 1.57 * BC - 1.30 * A^2 - 3.92 * B^2 - 4.22 * C^2
 \end{aligned}$$

B. 3: WL response surface plots of beetroot

Result represented in Fig. 3.7

X1=A: NaCl; X2=B: Sucrose; Actual factor C: Time = 4




Fig. 3.7: Effect of osmotic solution of NaCl, sucrose concentration on water loss during osmotic dehydration of beetroot cubes

A.4: Fitting models for RR of beetroot

The results presented in Table 3.13. The Model F-value of 3.26 implies the model is significant. There is only 3.99% chance that F-value will occur due to noise. Values of "Prob > F" less than 0.0500 indicate model terms are significant. In this case B, C are significant model terms. The "Lack of Fit F-value" of 16.02 implies that the Lack of Fit is significant. There is only 0.43% chance that a "Lack of Fit F-value" may occur due to noise.

Table – 3.13: ANOVA of response surface quadratic model for rehydration ratio

Analysis of variance table [Partial sum of squares - Type III]						
Source	Sum of Squares	df	Mean Square	F Value	p-value Prob > F	
Model	0.97	9	0.11	3.26	0.0399	significant
A-NaCl	0.025	1	0.025	0.76	0.4044	
B-Sucrose	0.54	1	0.54	16.23	0.0024	
C-Time	0.17	1	0.17	5.08	0.0478	
AB	1.404E-003	1	1.404E-003	0.042	0.8408	
AC	0.13	1	0.13	3.94	0.0754	
BC	6.272E-003	1	6.272E-003	0.19	0.6723	
A ²	0.033	1	0.033	1.00	0.3404	
B ²	6.902E-003	1	6.902E-003	0.21	0.6574	
C ²	0.053	1	0.053	1.59	0.2357	
Residual	0.33	10	0.033			
Lack of Fit	0.31	5	0.062	16.02	0.0043	significant
Pure Error	0.019	5	3.884E-003			
Total	1.30	19				

The final equation in terms of coded factors:

$$\begin{aligned}
 \text{RR-2} = & +2.71 + 0.043 * A - 0.20 * B - 0.11 * C - 0.013 * AB + 0.13 \\
 & * AC - 0.028 * BC + 0.048 * A^2 - 0.022 * B^2 - 0.060 * C^2
 \end{aligned}$$

B. 4: RR response surface plots of beetroot

Result represented in Fig. 3.8

X1=A: NaCl; X2=B: Sucrose; Actual factor C: Time = 4




Fig. 3.8: Effect of osmotic solution of NaCl, sucrose concentration on rehydration ratio during osmotic dehydration of beetroot cubes

The investigated data used for the optimization study were obtained using a Box Wilson *et al.* fractional factorial design (3 level-4 parameter), 20 runs. RSM was effective in optimizing the process of WR, WL, SG and RR by superimposition of the contour plots of all responses. Second-order polynomial (SOP) models for all system responses were statistically analyzed and found significant. Predicted and observed responses of carrot cubes in 50 °Brix of sucrose and 8% w/v NaCl for 4 hours were more significant at osmotic dehydration to achieve WR of 37.34%, SG of 14.33, WL of 51.34, and RR to be 3.176. While in case of beetroot 50°Brix of sucrose + 14% w/v NaCl for 4 hours were more significant at OD to achieve WR of 35.13, SG of 12.19, WL of 46.6, RR to be 3.031 as shown in the figure and table presented earlier, graphical techniques in association with RSM, aided in locating optimum operating conditions, which were experimentally verified and proven to be adequately reproducible.

3.4 SENSORY EVALUATION:

The results of sensory evaluation was represented at Table 3.14 and discussed below.

Table 3.14 Sensory evaluation of different food products.

Samples (Carrot & Beetroot Cubes)	Color and Appearance	Texture	Flavor	Taste	Overall Acceptability
Fresh	9.0	8.0	8.4	8.2	7.8
Salt treated	7.6	7.4	7.8	7.2	7.2
Sucrose treated	7.4	7.6	8.0	7.8	7.4
Salt + Sucrose treated	9.0	8.2	8.2	8.4	8.0

1) Color and Appearance:

Based on the data obtained from sensory evaluation by a panel of 6 member's color of health food formulations were found as 9.0, 7.6, 7.4 and 9.0 respectively on a hedonic scale. The carrot & beetroot cubes have dark orange & blackish-purple respectively, which is more attractive. It was evident that the color was significantly influenced by the difference in formulations of health food.

2) Texture:

Textures of food product formulations were found as 8.0, 7.4, 7.6 and 8.2 respectively. The texture of the product is free-flowing with very slight sickness.

3) Flavor:

It was found that the flavors of the food product formulations were found as 8.4, 7.8, 8.0 and 8.4 respectively. The formulations have a very good flavor.

4) Taste:

It was found that the taste of the Health food formulations was found as 8.2, 7.2, 7.6 and 8.0 respectively. The formulations have a very good taste except the salt-treated.

5) Overall acceptability

It was found that combination of salt + sucrose treated carrot & beetroot cubes were overall more acceptable as compared to salt and sucrose treated carrot and beetroot cubes.

CONCLUSION

Response Surface Methodology was the most effective for optimizing process parameters for osmotic dehydration of carrot and beetroot cubes in osmotic aqueous solution of sucrose and sodium chloride mixtures in a ratio of 1:5. The recommended process variables were 500 Brix of sucrose + 8 % w/v aqueous solution of sodium chloride for carrot and 500 Brix of sucrose + 14 % w/v NaCl solution for beetroot at osmotic dehydration for 4 hours considered to get maximum water loss, weight reduction, subsequent rehydration ratio, overall acceptability and minimum solute gain of rehydrated product. RSM was found to be effective for retaining flavor, odour and enhancing nutritional content. It can be concluded that osmotic dehydration had a positive impact on carrot & beetroot, retaining its nutritional value, flavor, odour, shelf life and preventing spoilage from microbial contamination. Osmotically dehydrated carrot and beetroot cubes can be used in noodles, soup, stews, pickles and casseroles.

Acknowledgement

The part of the work is financially supported by All India Council for Technical Education in the form of TE-QUIP fellowship provided to Mr. Shamsher Ali.

Abbreviations: RSM, Response Surface Methodology; WR, Weight Reduction; SG, Solute Gain; WL, Water Loss; RR, Rehydration Ratio; O.D., Osmotic Dehydration; CCD, Central Composite Design; RT, Room Temperature

REFERENCES

- [1] Akbarian, M., Ghasemkhani, N., & Moayedi, F. (2014). Osmotic dehydration of fruits in food industrial: A review. *Int. J. Biosci*, 4(1), 42-57. [View Article](#)
- [2] Ara, N., & Nur, H. (2009). In vitro antioxidant activity of methanolic leaves and flowers extracts of *Lippia alba*. *Research journal of medicine and medical sciences*, 4(1), 107-110. PMID:19579925 [View Article](#) [PubMed/NCBI](#)
- [3] Balaji, M., & Arshinder, K. (2016). Modeling the causes of food wastage in Indian perishable food supply chain. *Resources, Conservation and Recycling*, 114, 153-167. [View Article](#)
- [4] Bahmani, A., Jafari, S. M., Shahidi, S. A., & Dehnad, D. (2016). Mass transfer kinetics of eggplant during osmotic dehydration by neural networks. *Journal of food processing and preservation*, 40(5), 815-827. [View Article](#)
- [5] Box, G. E., & Wilson, K. B. (1951). On the experimental attainment of optimum conditions. *Journal of the royal statistical society: Series b (Methodological)*, 13(1), 1-38. [View Article](#)
- [6] Box, G. E., & Draper, N. R. (1987). *Empirical model-building and response surfaces* (Vol. 424). New York: Wiley.
- [7] Clifford, T., Bell, O., West, D. J., Howatson, G., & Stevenson, E. J. (2017). Antioxidant-rich beetroot juice does not adversely affect acute neuromuscular adaptation following eccentric exercise. *Journal of sports sciences*, 35(8), 812-819. PMID:27267689 [View Article](#) [PubMed/NCBI](#)
- [8] Di Pietro, N., Di Tomo, P., & Pandolfi, A. (2016). Carotenoids in cardiovascular disease prevention. *JSM Atheroscler*, 1(1), 1002.
- [9] Dolker, D., Bakshi, P., Wali, V. K., Dorjey, S., Kour, K., & Jasrotia, A. (2017). Integrated Nutrient Management in Fruit Production. *Int. J. Curr. Microbiol. App. Sci*, 6(7), 32-40. [View Article](#)
- [10] Gupta, S., Singh, K., & Gupta, V. (2019). Genetic resource conservation of horticultural crops in India-Achievements and issues. *Progressive Horticulture*, 51(1), 16-29. [View Article](#)
- [11] Gupta, N., & Shukla, R. N. (2017). Preparation and Quality Evaluation of Dehydrated Carrot and Onion Slices. *Journal of Food Processing & Technology*. [View Article](#)
- [12] Gonçalves, E. M., Pinheiro, J., Abreu, M., Brandão, T. R., & Silva, C. L. (2010). Carrot (*Daucus carota* L.) peroxidase inactivation, phenolic content and physical changes kinetics due to blanching. *Journal of Food Engineering*, 97(4), 574-581. [View Article](#)
- [13] Handbook on Horticulture Statistics, 2018. Department of Agriculture, Ministry of Agriculture & Farmers' Welfare, Government of India. Government of India.

- [14]Kapadia, J G., A Azuine, M., Subba Rao, G., Arai, T., Iida, A., & Tokuda, H. (2011). Cytotoxic effect of the red beetroot (*Beta vulgaris* L.) extract compared to doxorubicin (Adriamycin) in the human prostate (PC-3) and breast (MCF-7) cancer cell lines. *Anti-Cancer Agents In Medicinal Chemistry (Formerly Current Medicinal Chemistry-Anti-Cancer Agents)*, 11(3), 280-284. PMID:21434853 [View Article](#) [PubMed/NCBI](#)
- [15]Kobaek-Larsen, M., Nielsen, D. S., Kot, W., Krych, Ł., Christensen, L. P., & Baatrup, G. (2018). Effect of the dietary polyacetylenes falcarinol and falcariindiol on the gut microbiota composition in a rat model of colorectal cancer. *BMC research notes*, 11(1), 411. PMID:29945666 [View Article](#) [PubMed/NCBI](#)
- [16]Madison, D., & Coleman, E. (2007). *Preserving Food Without Freezing Or Canning: Traditional Techniques Using Salt, Oil, Sugar, Alcohol, Vinegar, Drying, Cold Storage, and Lactic Fermentation*. Chelsea Green Publishing.
- [17]Mercali, G. D., Marczak, L. D. F., Tessaro, I. C., & Noreña, C. P. Z. (2011). Evaluation of water, sucrose and NaCl effective diffusivities during osmotic dehydration of banana (*Musa sapientum*, shum.). *LWT-Food Science and Technology*, 44 (1), 82-91. [View Article](#)
- [18]Mudahar G. S., Toledo R. T., Floros J. D. and Jen J. J. (1989). Optimization of carrot dehydration process using response surface methodology. *Journal of Food Science* Vol. 54, Pp 174-719. [View Article](#)
- [19]Pryor, B. M., Davis, R. M., & Gilbertson, R. L. (2000). A toothpick inoculation method for evaluating carrot cultivars for resistance to *Alternaria radicina*. *HortScience*, 35(6), 1099-1102. [View Article](#)
- [20]Rundla, A., Mishra, A. A., & Shukla, R. N. (2018). Studies on Physical and Organoleptic Properties of Osmotically Dehydrated Carrot (*Daucus carota* L.) Slices. *Int. J. Curr. Microbiol. App. Sci*, 7(9), 2457-2468. [View Article](#)
- [21]Singh, R. B., Rastogi, V., Singh, R., Niaz, M. A., Srivastav, S., Aslam, M., ... & Postiglione, A. (1996). Magnesium and antioxidant vitamin status and risk of complications of ageing in an elderly urban population. *Magnesium research*, 9 (4), 299-306.
- [22]Singh, D. P., Beloy, J., McInerney, J. K., & Day, L. (2012). Impact of boron, calcium and genetic factors on vitamin C, carotenoids, phenolic acids, anthocyanins and antioxidant capacity of carrots (*Daucus carota*). *Food chemistry*, 132(3), 1161-1170. PMID:29243596 [View Article](#) [PubMed/NCBI](#)
- [23]Singh, A., Verma, S., Singh, V., Nanjappa, C., Roopa, N., Raju, P., & Singh, S. N. (2015). Beetroot juice supplementation increases high density lipoprotein-cholesterol and reduces oxidative stress in physically active individuals. *J Pharm Nut Sci*, 5(3), 179-85. [View Article](#)
- [24]Shah, N. S., & Nath, N. (2008). Changes in qualities of minimally processed litchis: Effect of antibrowning agents, osmo-vacuum drying and moderate vacuum packaging. *LWT-Food Science and Technology*, 41(4), 660-668. [View Article](#)
- [25]Shi J. and Maguer M. L. (2002). Osmotic dehydration of foods, mass transfer and modelling aspects. *Food reviews international* Vol 18, 2002, Pp 305-335. [View Article](#)
- [26]Tan, K. W., Killeen, D. P., Li, Y., Paxton, J. W., Birch, N. P., & Scheepens, A. (2014). Dietary polyacetylenes of the falcarinol type are inhibitors of breast cancer resistance protein (BCRP/ABCG2). *European journal of pharmacology*, 723, 346-352. PMID:24269959 [View Article](#) [PubMed/NCBI](#)
- [27]Venter, G., Haftka, R., & Starnes, Jr, J. (1996). Construction of response surfaces for design optimization applications. In *6th Symposium on Multidisciplinary Analysis and Optimization* (p. 4040). [View Article](#)