

SDRP Journal of Aquaculture, Fisheries & Fish Science (ISSN: 2575-7571)

Trace element concentrations in sediments and water from Bahia de Lobos, Sonora, Mexico.

DOI: 10.25177/JAFFS.2.2.1

Research

Received Date: 26th Jul 2018Accepted Date: 01st Sep 2018Published Date: 07th Sep 2018

Copy rights: © This is an Open access article distributed under the terms of International License.



Solorzano-Meza Jose Luis^{1,2}, Meza-Montenegro Maria Mercedes^{1*}, Mondaca-Fernandez Iram¹, Gortares-Moroyoqui Pablo¹, Vega-Millan Christian Beatriz^{1,2}, Balderas-Cortes Jose de Jesus^{1,*}.

¹Instituto Tecnológico de Sonora. 5 de Febrero 818 sur, Cd. Obregon, Sonora, Mexico. 85000.

²Programa de Doctorado en Ciencias Especialidad en Biotecnología. Departamento de Biotecnología y Ciencias Alimentarias, Instituto Tecnológico de Sonora. 5 de Febrero 818 sur, Cd. Obregon, Sonora, Mexico. 85000.

*These two authors contributed equally with the manuscript.

CORRESPONDENCE AUTHOR

Jose de Jesus Balderas Cortes

Tel: +52 1 (644) 410 90 00; Email: jose.balderas@itson.edu.mx;

CITATION

Jose Luis Antonio Solorzano Meza, Trace element concentrations in sediments and water from Bahia de Lobos, Sonora, Mexico. (2018) SDRP Journal of Aquaculture, Fisheries & Fish Science 2(2)

ABSTRACT

Superficial water and sediment samples were collected from Bahia de Lobos in Sonora. This is the first work to describe the analysis of heavy metals in water of Bahia de Lobos including the measurement of Arsenic and Mercury. We determined Lead (Pb), Cadmium (Cd), Copper (Cu), Arsenic (As) and Mercury concentrations. This estuary was sampled monthly from May to December of 2016, and samples were analyzed by Atomic Absorption Spectrometry (AAS) and AAS coupled to Hydride Generation. The levels of metals found in water were: Pb (1080 $\mu\text{g L}^{-1}$), Cu (104 $\mu\text{g L}^{-1}$), Cd (80 $\mu\text{g L}^{-1}$), As (<LOQ: Limit Of Quantification), Hg (<LOQ). Lead exceeded the Mexican regulatory limits (200 $\mu\text{g L}^{-1}$) for Protection of Aquatic Life (PAL). Cd exceeded the established values (3.9 and 1.0 $\mu\text{g L}^{-1}$) by the USEPA for the Protection of Aquatic Life Fresh Acute Criteria (PALFAC) and the Canadian Council of Ministers of the Environment (CCME) respectively, as well as the level permitted by the Protection of Aquatic Life Marine Acute Criteria (PALMAC)

(0.12 $\mu\text{g L}^{-1}$). Copper (104 $\mu\text{g L}^{-1}$) was above the USEPA in PALFAC (18 $\mu\text{g L}^{-1}$). Levels of Pb (1.85 mg kg^{-1}), Cd (<LOQ), Cu (4.52 mg kg^{-1}), As (1045 $\mu\text{g kg}^{-1}$), and Hg (2.76 $\mu\text{g kg}^{-1}$) were also found in sediments. Pb exceeded the International Legislation according to the Screening Quick Reference Tables (US-SQUIRTS) established for mammals (0.0537 mg kg^{-1}) and Hg exceeded PALFAC (0.17 $\mu\text{g kg}^{-1}$), PALMAC (0.13 $\mu\text{g kg}^{-1}$), and SQUIRTS for invertebrates and plants (0.1 and 0.3 $\mu\text{g kg}^{-1}$). Only temporal differences for Pb in water ($P=0.0270$), and for Cu in sediment were found ($P=0.0000$), while As presented spatial temporal differences in water and sediment, respectively. Consequently, the metal levels found in water and sediment from Bahia de Lobos might cause potential adverse effects in the biota and it is necessary to work with the Mexican legislation to prevent further contamination in the Bahia.

Keywords: trace elements, sediment, water, Bahia de Lobos, pollution

INTRODUCTION

Estuaries are considered to be a sink for chemical substances which can adhere to particulates suspended in water and then deposit in sediments, repeating the process cyclically [1-2]. Estuaries and coastal areas exhibit human caused pollution that compromises their ecological integrity [3-6].

The bays provide biological protection for endemic species and serve as breeding and nesting sites as well. Bays are the final recipients of human pollution but also provide food for humans and trace elements can return into the human diet by biomagnification, through the trophic chain [7].

Trace elements occur naturally in the environment, but various human activities such as mining, industry, and agriculture increase their concentration [8-10].

Pb and Cd tend to accumulate in the bone structure of organisms due to their chemical similarity to calcium, causing a decrease in the quality of life of marine organisms at low exposure levels. In addition, they cause chronic adverse effects in the reproductive system [11-17]. Arsenic is carcinogenic and considered the most toxic element, damaging the nervous and vascular systems [18-19]. Hg has the ability to move from the sediment into the water and as it is lipophilic, is bioaccumulated in the reproductive and nervous systems, mainly in the brain [20-21]. Cu regulates some natural functions in the marine biota working like an enzyme necessary for living organisms [22]. However, it has been reported that at concentrations between 10-15 times higher than the required levels, it produces adverse effects in aquatic organisms [23].

The importance of monitoring metal levels in sediment is that the metals adhere to the particulate matter, accumulating over time and are re-suspended into water with the tidal cycles, and become bioavailable to organisms in the marine environment [24-26].

Recent studies carried out in a variety of countries (Venezuela, Argentina, Canada, Tunisia, Pakistan, Egypt, France, China and Russia) have shown that heavy metals are chemically very stable, i.e. they do not degrade over time. On the contrary, their presence increases over time in sediment, water, soil, air and

biota [27--35].

In Mexico, very few studies of heavy metals in water and sediments from estuaries have been reported since 1980. Cadmium (0.60 mg L^{-1}) was found in water from Laguna Madre Tamaulipas by Pulich (1980) [36]. González-Lozano et al. (2006) [37] reported Cu (2.06 mg kg^{-1}), Pb (3.3 mg kg^{-1}) and Cd (937.5 mg kg^{-1}) in sediment at Salina Cruz, Oaxaca. Celis-Hernández et al. (2017) [38] determined Pb (17 mg kg^{-1}), Cu (25 mg kg^{-1}) and As ($14\,000 \text{ } \mu\text{g kg}^{-1}$) were present in sediment from Veracruz, and Botello et al. (2018) [39] detected concentrations of Pb (40.35 mg kg^{-1}), Cd ($0.0417 \text{ mg kg}^{-1}$), Cu (52.59 mg kg^{-1}) and Hg ($210 \text{ } \mu\text{g kg}^{-1}$) in sediment, which represents the contamination from these elements for the last 40 years.

Sonora is located in northwest Mexico and borders the United States to the north. It is located in the Pacific basin and is bounded on the west by the Sea of Cortez [40]. Bahia de Lobos is located in southern Sonora on the eastern coast of the Gulf of California and is the main commercial source for fish in Sonora. Additionally, this bay is the final discharge area of the three major waste water channels (agricultural, domestic and industrial) that come from the city of Obregon and from the primary Yaqui communities. Ciudad Obregon and the Yaqui communities treat only a 9.5% of the wastewater Gortares-Moroyooqui, (2010) [41]. As a consequence, the biota from this bay is in constant environmental stress from the introduction of various pollutants. Bahia de Lobos has four types of mangroves, species protected by Mexican Legislation which are of global interest for their key role in the reproduction cycle of some marine species, e.g. shrimp [42].

Studies of the presence of metals in this area are scarce. One of the most recent was conducted by Vargas-Gonzalez et al. (2017) [43] who found Pb, Cd and Cu in sediments of Bahia de Lobos. The reported concentrations were 18.5 mg kg^{-1} , 1.22 mg kg^{-1} and 16.30 mg kg^{-1} respectively. Ortega and Vasquez (1992) [44] carried out a study of sediment from the Bahia de Lobos, quantifying Pb and Cu levels, and found 59 mg kg^{-1} and 88 mg kg^{-1} respectively. García-Rico et al. (2004) [45] analyzed metals in oyster cul-

ture areas, in the Northern and Central coasts of Sonora, quantifying heavy metals in sediments, and detecting the highest concentrations in Guaymas, Sonora (Cu 14.85 mg kg⁻¹; Cd 5.62 mg kg⁻¹; and Pb 46.55 mg kg⁻¹).

Due to the scarcity of studies of the negative impact that these metals are causing in the biota of the Bahía de Lobos, the goal of this study was to analyze the concentration of the most toxic trace elements, As, Hg, Pb, Cd and Cu in sediments including for the first time determination of levels of arsenic and mercury, and the analysis of the study metals in water from bahía de Lobos.

MATERIALS & METHODS

Location of the study site

Bahía de Lobos is located in south Sonora, Mexico (Figure 1), and has an area of 11, 978 ha. The estuary is considered of great biological importance because it has four mangrove species at risk of extinction according to the Official Mexican Norm [46], and serves as a safe place for reproduction and nesting of endemic and step species. Bahía de Lobos has an average annual fishery production of \$37,500 per hectare of mangrove [42].




Figure 1. Sampling sites in Bahía de Lobos, Sonora.

Sampling sites

Five points (latitude and longitude P1:27° 21'7.59"N,110°27'17.68"O; P2: 27°20'40.75"N, 110° 28'2.80"O; P3: 27°20'9.99"N, 110°28'54.97"O; P4: 27° 18'8.18"N, 110°29'16.96"O; P5: 27°18'14.66"N, 110° 30'59.31"O) of the bay were selected for monthly sampling based on the estuarine area (Fig. 1). The collection of sediments and water samples was done from May to December, 2016. The points were strategically chosen and marked with a GPS Garmin, model ETrex 10, to establish the temporal space distribution from the point of discharge.

Collection of sediments

The top 10 cm of surface sediments were collected using a plastic shovel and placed in a 500 mL polypropylene flask. Samples were transported in coolers at 4 °C to the Laboratory of Toxicology and Public Health at the Instituto Tecnológico de Sonora. The sediment was dried at 45 °C in a Model 107800 Boekel Scientific Oven and then ground with a mortar and pestle and packed in polyethylene bags at -20 °C until the analysis could be completed.

Collection of water

The surface water samples were collected using a 500 mL polypropylene flask and transported in coolers at 4 °C to the Laboratory of Toxicology and Public Health at the Instituto Tecnológico de Sonora, using protocol established by Mexican Legislation [47].

Samples processing and analysis

All glassware used for the metal analysis was previously soaked in 10% HNO₃ (v/v) and rinsed with deionized water.

Water and Sediment

For sediment chemical analyses, five grams of dried sediment were placed in a 250 mL Pyrex volumetric flask, a total of 20 mL of concentrated HNO₃ grade Omnitrace was added to the flask and placed on a heating plate for digestion according to the Mexican Norm [48]. Then 5 mL of nitric acid were added intermittently to digest the sediment completely until the supernatant became amber-clear. When the sample was nearly dried, the solution was filtered through a 0.42 µm membrane into a 100 mL Pyrex volumetric flask and then filled with deionized water.

For water chemical analyses, 30 mL of water were placed into a 250 mL Pyrex volumetric flask, a total of 20 mL of concentrated HNO₃ grade Omnitrace was added to the flask and placed on a heating plate for digestion according the Mexican Norm [47]. Then 5 mL of nitric acid were added again to digest the water until the supernatant became amber-clear. After the sample was nearly dried, the solution was filtered through a 0.42 µm membrane in a 100 mL Pyrex volumetric flask and filled with deionized water.

After sample digestion, Pb, Cd, Hg and Cu were determined to be present by Atomic Absorption Spectrometry (AAS). As and Hg were analyzed using Hydride Generator (HG) coupled to AAS equipment. The presence of Arsenic was determined by flame and Hg by cold vapor according to the Mexican Norm [47-48]. For quality control the duplicate samples were analyzed and three blanks and three spikes were prepared for every 10 samples analyzed. The sediment recoveries for the metals ranged from 95.1

to 106.79%, and the variation coefficients (CV) values for the metals Cu, Cd, Pb, As and Hg were 0.42, 0.46, 27.13, 8.51 and 6.58%, respectively.

The water recoveries for heavy metals ranged from 90.75 to 104.25%, and the CV values for Cu, Cd, Pb, As, and Hg were 0.28, 1.59, 9.60, 3.07 and 4.23%, respectively.

Statistical analyses

Measures of central tendency of the levels of metals in the water and sediment were obtained. To compare the metal concentrations with the sampling points and months, a factorial of 95% ANOVA was used. Later, in the cases in which a statistically significant difference presented, a media difference analysis (Fisher LSD) was carried out using STATGRAPHICS Plus software for Windows 5.1 [50].

RESULTS & DISCUSSION

The mean levels of metals in the water samples from Bahia de Lobos, Sonora, are shown in Table 1. The mean concentration of Pb was 1080 µg L⁻¹, 2160 times higher than the value reported in Kola, Russia (0.5 µg L⁻¹) by Moiseenko et al. (2018) [35], and 13.5 times higher than the levels reported in Cordoba, Argentina (186.3 µg L⁻¹) by Griboff et al. (2018) [30].

The average level of Cd in the water was 80 µg L⁻¹, 266 times higher than that found in Kola, Russia (0.3 µg L⁻¹) by Moiseenko et al. (2018) [35]. The average level of Cu detected was 104 µg L⁻¹, 8.5 times higher than values found in Kola, Russia (12.1 µg L⁻¹) by Moiseenko et al. (2018) [35], and 12.5 times lower than reported in Cordoba, Argentina (1300 µg L⁻¹) by Griboff et al. (2018) [30]. The concentrations of As and Hg in the water were below the limit of detection. There are few studies which report levels of these elements in water despite their important toxicity. Griboff et al. (2018) [30] found levels of arsenic at 1470 µg L⁻¹ in Cordoba, Argentina and Sprague and Vermaire (2018) [27] detected 972 µg L⁻¹ in Ontario, Canada. Perez-Zapata (1981) [53] reported 4 µg L⁻¹ of Mercury, in Veracruz, Mexico. In our study, the average levels of Pb in the water were 1080 µg L⁻¹. The average levels of Cd were 80 µg L⁻¹,

which did not exceed the Mexican Norm [49] ($100 \mu\text{g L}^{-1}$) for Estuaries and Protection of Aquatic Life (PAL), but did exceed by 20.5 times the maximum established criteria ($3.9 \mu\text{g L}^{-1}$) for the Protection of Aquatic Life, Fresh Acute Criteria (PALFAC) from United States Environmental Protection Agency (USEPA) [52]. The Cd levels exceeded the value established by the Protection of Aquatic Life and Fresh Acute Criteria (PALFAC) from Canadian Council of Ministers of the Environment (CCME)

[53] by 80 times and were 666 times above the value established in the Protection of Aquatic Life and Marine Acute Criteria (PALMAC) from CCME [53] (Table 1). The Cu levels in the water were $104 \mu\text{g L}^{-1}$ and did not exceed the Mexican Legislation for estuaries nor the Mexican Guideline for the Protection of Aquatic Life (PAL) [49], but the value established by the USEPA-PALFAC [52] ($18 \mu\text{g L}^{-1}$) was 5.7 times higher in the Bahia de Lobos (Table 1).

Table 1. Comparison of mean metal levels in water from Bahia de Lobos, Sonora, Mexico, and other studies carried out around the world with the International Norms or Criteria for the protection of aquatic life.

Country	Studied elements ($\mu\text{g L}^{-1}$)					References	Norms/ Criteria
	Pb	Cd	Cu	As	Hg		
Meiliang Bay, China	6000	740	1060	--	--	Rajeshkumar et al. (2015)	
Kabul, Pakistan	186.3	90.6	32	--	--	Khan et al. (2018)	
Kola, Russia	0.5	0.30	12.1	--	--	Moiseenko et al. (2018)	
Ontario, Canada	--	--	--	972	--	Sprague and Vermaire, (2018)	
Cordoba, Argentina	80	<LOD	1300	1470	LOD	Griboff et al. (2018)	
Veracruz, México	430	--	--	--	4	Perez-Zapata, (1981)	
Bahia de Lobos	1080+1370	80+205	104+61.2	<LOQ	<LOQ	Our study	
Mexico	200	100	4000	100	5	NOM-001-ECOL-1996	PAL
Mexico	200	100	4000	100	10	NOM-001-ECOL-1996	Estuaries
USA	-/-	-/-	-/-	69	2.1	EPA	PALMAC
USA	-/-	3.9	18	360	2.4	EPA	PALFAC
Canada	-/-	1	-/-	5	0.026	CCME	PALFAC
Canada	-/-	0.12	-/-	12.5	0.016	CCME	PALMAC

-/- Without information; -- No analyzed; LOD: Limit of detection; LOD-Hg: $1.27 \mu\text{g/L}$; LOD-As: $0.92 \mu\text{g/L}$; LOQ: Limit of quantification; LOQ-Hg: $2.09 \mu\text{g/L}$; LOQ-As: $3.3 \mu\text{g/L}$; PAL: Protection of the aquatic life; PALMAC: Protection of the Aquatic Life, Marine Acute Criteria; PALFAC: Protection of the Aquatic Life, Fresh Acute Criteria; CCME: Canadian Council of Ministers of the Environment.

The average values for metals in sediment samples from the Bahía de Lobos are shown in Table 2. The mean concentration of Pb in sediments was 1.85 mg kg^{-1} , a value 4.7 times lower than the value reported in Cordoba, Argentina by Griboff et al. (2018) [30], [Table 2], and also 2.4 times lower than the levels reported in Lake Bourget, France (4.47 mg kg^{-1}) by Lécivain et al. (2018) [Table 2]. The average concentrations of Cd were below the limit of quantification (0.08 mg kg^{-1}), but levels of Cadmium in the sediment have been reported in Cordoba, Argentina ($0.0305 \text{ mg kg}^{-1}$) and Lake Bourget, France (0.09 mg kg^{-1}) by Griboff et al. (2018) and Lécivain et al. (2018) respectively (Table 2) [30, 33]. The average concentrations of Cu in the sediment from Bahía de Lobos were 4.52 mg kg^{-1} , 1.17 times higher than that found in Lake Bourget, France (3.86 mg kg^{-1}) by Lécivain et al. (2018) and similar to those reported in Cordoba, Argentina (5.6 mg kg^{-1}) by Griboff et al. (2018) [Table 2]. The mean concentrations of As in sediment from Bahía de Lobos were $1045 \text{ } \mu\text{g kg}^{-1}$. Arsenic has not been monitored in most of the recent works, except in Argentina where a value of $1900 \text{ } \mu\text{g kg}^{-1}$ was reported [33]. This latter value was 1.8 times higher than the value found in this study. Mercury had an average level of $2.76 \text{ } \mu\text{g kg}^{-1}$ in the sediment from Bahía de Lobos (Table 2). There are unfortunately no reports of the analysis of this metal in sediment in recent investigations, even though this element is considered bioaccumulative and neurotoxic, capable of causing serious damage to aquatic life. Only Griboff et al. (2018) [30] reported levels of $0.052 \text{ } \mu\text{g kg}^{-1}$ of Hg in sediment from Cordoba, Argentina, a value 53 times lower than was found in this study (Table 2).

The concentration of Pb in the sediment from Bahía de Lobos was 1.85 mg kg^{-1} . In the absence of a Mexican regulation, this result was compared to International Guidelines, and was found to exceed by 34.4 times the value of $0.0537 \text{ mg kg}^{-1}$ established by Buchman (2008) [54] as the threshold necessary for the protection of mammals (Table 2). Cd concentrations in sediment from Bahía de Lobos were below our limit of detection. The concentration of Cu was 4.52 mg kg^{-1} , and did not exceed any International

Norm from USA and Canada, but this value was close (5.4 mg kg^{-1}) to the levels in Screening Quick Reference Tables (SQUIRTS) established for the protection of mammals (Buchman, 2008) [54]. The As concentration in the sediment of Bahía de Lobos was $1045 \text{ } \mu\text{g kg}^{-1}$, and did not surpass the value established for the Protection of the Aquatic Life by the Norms of the USA and Canada.

The average concentration of Hg in sediment of this study was $2.76 \text{ } \mu\text{g kg}^{-1}$, and exceeded by 21 times the established value ($0.13 \text{ } \mu\text{g kg}^{-1}$) by the Protection of Marine Aquatic Life (PMAL) of the Canadian Council of Ministers of the Environment (CCME) [53] and by 16 times the concentration established ($0.17 \text{ } \mu\text{g kg}^{-1}$) by Protection of Fresh Aquatic Life (PFAL) of CCME [53]. The concentration in the Bahía de Lobos was 27.6 times above the levels established ($0.1 \text{ } \mu\text{g kg}^{-1}$) by Buchman (2008) [54] in the SQUIRTS and 9.2 times the established value ($0.3 \text{ } \mu\text{g kg}^{-1}$) in the SQUIRTS for plants.

The averages of metal concentrations in the sediments were higher than in the water, with the exception of the Cd which was higher in water. That is not surprising since these contaminants are found in higher concentrations adhered to the sediments and, with the tides, are re-suspended cyclically in the water. The behavior observed in this study is similar to that reported in the literature [1-2].

The metals present in the water exceed the Mexican regulations for the Protection of Aquatic Life, but in Mexico regulations for estuarine sediments do not exist. In the same way, the thresholds of the USEPA [52], CCME [53], and SQUIRTS [54] were surpassed. In these conditions, the marine aquatic life and freshwater species have been compromised because both marine and freshwater organisms live in the estuaries. These results may be due to the variety of contamination sources, for example, the lead present in gasoline and some pesticides in the 90's, often present in automobile parts and paint, which wear out and leach into the environment, contamination from industry, and household waste generation. Some of these contaminants are transported and discharged into the Bahía de Lobos from the three major waste water channels which come from Obregon and some

Yaqui communities. In addition, the levels of metals established by Buchman (2008) [54] in the SQUIRTS for mammals, invertebrates and plants were surpassed. These results showed that the trace metals found in Bahia de Lobos could generate adverse effects to living organisms.

Table 2. Comparison of mean levels of metals in sediment from Bahia de Lobos, Sonora, Mexico with the Mexican and International Norms for the Protection of Aquatic Life.

Country	Determined elements (mg kg ⁻¹)					References	Norms/Criteria
	Pb	Cd	Cu	As*	Hg*		
Meiliang Bay, China	8.53	0.50	7.19	--	--	Rajeshkumar et al. (2017)	
Kabul, Pakistan	31.9	7.1	3.6	--	--	Khan et al. (2018)	
Lake Bourget, France	4.47	0.09	3.86	--	--	Lécrivain et al. (2018)	
Sfax, Tunisia	68.6	11.1	34.3	--	--	Bahloul et al. (2018)	
Red sea, Egypt	75	3.83	108	--	--	El-Taher et al. (2017)	
Cordoba, Argentina	8.7	0.0305	5.6	1900	0.052	Griboff et al. (2018)	
Mar Caribe, Venezuela	<LOD	<LOD	0.5	--	--	Urbina-Barreto et al. (2014)	
Mexico	17	--	25	14000	--	Celis-Hernandez et al. (2017)	
Bahia de Lobos	1.85±5.98	<LOQ	4.52±7.04	1045±1226	2.76±10.34	Our study	
Canada	35	0.7	35.7	7240	0.17	CCME	PFAL, ISQG
Canada	30.2	0.6	18.7	5900	0.13	CCME	PMAL, ISQG
USA	11	0.77	28	43000	-/-	Buchman, (2008)	SQUIRTS, Avian
USA	500	20	50	60000	0.1	Buchman, (2008)	SQUIRTS, Inverts
USA	0.0537	0.0022	5.4	5700	-/-	Buchman, (2008)	SQUIRTS, Mammals
USA	50	4	70	18000	0.3	Buchman, (2008)	SQUIRTS, Plants

*Concentration is in µg kg⁻¹; -/- Without information; -- Not analyzed; LOD: Limit of detection; LOD-Cd: 0.02 mg/kg; LOQ: Limit of quantitation; LOQ-Cd: 0.08 mg/kg; ISQG: Interim sediment quality guidelines; PFAL: Protection of fresh aquatic life; PMAL: Protection of marine aquatic life; SQUIRTS: Screening quick reference tables; CCME: Canadian Council of Ministers of the Environment.

The spatial temporal variation of metal levels in water and sediment from Bahia de Lobos, is shown in Table 3. For water, statistically significant differences of Pb levels are presented by month ($P= 0.0270$), but not by sampling point. In July, the higher levels (2.54 mg L^{-1}) were found and these values were statistically different from May (0.47 mg L^{-1}), June (0.47 mg L^{-1}), October (0.47 mg L^{-1}), November (0.47 mg L^{-1}) and December (0.47 mg L^{-1}), but statistically equal to August (1.709 mg L^{-1}) and September (1.835 mg L^{-1}). On the other hand, for Cd, Cu, As and Hg, statistically significant differences in their spatial (sampling sites) and temporal (sampled months) concentrations were not found. In the sediment, Cu presented statistically significant temporal differences in its levels ($P= 0.0000$). Table 3, shows higher concentrations (18.93 mg kg^{-1}) for July, and these levels were statistically different from May (0.007 mg kg^{-1}), June (0.007 mg kg^{-1}), August (4.25 mg kg^{-1}), September (2.21 mg kg^{-1}), October (2.16 mg kg^{-1}), November (2.48 mg kg^{-1}) and December (5.23 mg kg^{-1}). For Pb,

Cd, and Hg, statistically significant differences in temporal concentrations were not found. Arsenic presented a temporal statistically significant difference ($P= 0.0119$), as well as a spatial variation in its levels between the five sampling points: 3 (2.316 mg kg^{-1}), 1 (0.555 mg kg^{-1}), 2 (0.822 mg kg^{-1}), 4 (0.885 mg kg^{-1}) and 5 ($0.0059 \text{ mg kg}^{-1}$), respectively. The highest levels of Cu and Pb were detected in July (Table 3). July saw the heaviest precipitation in Sonora for the entire year as measured by the National Commission of Water [55]. This could account for the high levels of metals, because of possible resuspension of sediments which would change the metal levels in the superficial water and sediment. For the same reason we detected the highest levels of Pb in December (sediments) because October to December low precipitation was recorded [55] and this influences the concentration of the metals in sediments. And the high Cu sediment levels in July can be affected by heavier precipitations removing the metals in sediments. Some can be detected in water and others has more affinity to particulate matter like Cu.

Table 3. Means of metals concentrations in water and sediment from analyzed months of Bahia de Lobos.

Month	Sediment (mg kg^{-1})					Water (mg L^{-1})				
	Pb	Cd	Cu	As	Hg	Pb	Cd	Cu	As	Hg
May	0.47 ^a	0.02 ^a	0.007 ^a	0.78 ^a	0.001045 ^a	0.47 ^a	0.04 ^a	0.106 ^a	0.00046 ^a	0.00104 ^a
June	0.47 ^a	0.02 ^a	0.007 ^a	1.89 ^{bc}	0.001045 ^a	0.47 ^a	0.04 ^a	0.039 ^a	0.00046 ^a	0.00104 ^a
July	0.47 ^a	0.02 ^a	18.93 ^b	0.73 ^{ab}	0.001045 ^a	2.54 ^b	0.04 ^a	0.087 ^a	0.00046 ^a	0.00104 ^a
August	0.47 ^a	0.02 ^a	4.25 ^a	0.92 ^{ab}	0.001045 ^a	1.709 ^{ab}	0.04 ^a	0.083 ^a	0.00046 ^a	0.00104 ^a
September	0.47 ^a	0.02 ^a	2.21 ^a	0.276 ^a	0.001045 ^a	1.835 ^{ab}	0.029 ^a	0.102 ^a	0.00046 ^a	0.00104 ^a
October	6.46 ^a	0.02 ^a	2.16 ^a	0.958 ^{bc}	0.001045 ^a	0.47 ^a	0.04 ^a	0.162 ^a	0.00046 ^a	0.00104 ^a
November	0.47 ^a	0.02 ^a	2.48 ^a	2.239 ^c	0.001045 ^a	0.47 ^a	0.04 ^a	0.132 ^a	0.0110 ^a	0.00104 ^a
December	5.27 ^a	0.02 ^a	5.23 ^a	0.518 ^b	0.001045 ^a	0.47 ^a	0.04 ^a	0.13 ^a	0.00046 ^a	0.00104 ^a

^{abc}Means with different superscripts indicate a statistically significant difference ($P<0.050$) LSD test.

CONCLUSIONS

In spite of the fact that Cu does not exceed the limits allowed by Mexican regulations in water, it does exceed those established by the US-EPA and CCME. This means that Mexican regulations lag behind and expose deficiencies in the environmental care policies. In addition, the treatment plants in the area only treat 9.5% of the wastewater that reaches the site. This legislative omission is responsible for contaminants reaching the Bay which may be causing these elements to accumulate in the bone structure of the organisms and cause damage to their nervous systems.

The Mexican legislation does not have a norm for heavy metals in sediments and thus, limits protection for the biota.

The USA and Canadian norms were useful to fill the legislative gap in this work.

Pb exceeds the protective threshold for mammals in SQUIRTS and thus can produce problems in their reproductive systems.

Hg surpasses the limits set to protect marine aquatic life established by CCME and SQUIRTS for plants. This element is lipophilic and has affinity to nervous systems and can affect the cerebral synapses. It also exceeded the limit established for plants. It must be remembered that in Bahía de Lobos there are four types of mangroves, protected by national and international standards because of their great importance for the environment. This is because they are ecological niches of a great variety of marine species, and barriers that prevent pollution from reaching the sea in greater quantity. They are also generators of wealth in local and national economies.

Continued monitoring of this bay is recommended as well as conducting a more exhaustive study including the endemic biota, to enable remedial actions if necessary.

ACKNOWLEDGEMENTS

The authors would like to thank Instituto Tecnológico de Sonora for financing this study with the following generous grants: PROFAPI_CA_2018-1145 and

PROFAPI_2018_0030. In addition, thanks to Dr. Paul W. Kilpatrick for his support with the English edition. The publication of this manuscript was made possible with the support of the Program for Strengthening Educational Quality (PFCE), 2018.

REFERENCES

- [1] Turner, A., Le Roux, S.M., Millward, G.E. (2004). Speciation and partitions of cadmium and zinc in two contrasting estuaries: The role of hydrophobic organic matter. *Limnology and Oceanography*, 49, 11-19. [View Article](#)
- [2] Segura, R., Arancibia, V., Zú-iga, M.C., Pastén, P. (2006). Distribution of copper, zinc, lead and cadmium concentrations in stream sediments from the Mapocho River in Santiago, Chile. *Journal of Geochemical Exploration*, 91, 71–80. [View Article](#)
- [3] Kulikova, I., Seisum, Z., Legzdina, M. (1985). Heavy metals in marine organisms. *SympBiolHung*, 29, 141–151.
- [4] Nelson, G.C. (2005). Drivers of ecosystem change: Summary chapter. In *Island press* (Ed). *Ecosystems and human well-being* (pp. 73-76). Washington, USA.
- [5] Hughes, T.P., Bellwood, D.R., Folke, C., Steeneck, R.S., Wilson, J. (2005). New paradigms for supporting resilience in marine ecosystems. *Trends in Ecology and Evolution*, 20(7), 380-385. PMID:16701400 [View Article](#) [PubMed/NCBI](#)
- [6] Ghosh, U., Luthy, R.G., Cornelissen, G., Werner, D., Menzie, C. (2011). In-situ sorbenamendments: a new direction in contaminated sediment management. *Environmental Science and Technology* 45(4), 1163-1168. PMID:21247210 PMID:PMC3037809 [View Article](#) [PubMed/NCBI](#)
- [7] Roberts, L., Brower, A. L., Kerr, G. N., et al. (2015). The nature of wellbeing: how nature's ecosystem services contribute to the wellbeing of New Zealand and New Zealanders. Department of Conservation (Ed).
- [8] Zarazúa, G., Tejeda, S., Ávila, P., et al. (2011). Metal content and elemental composition of particles in cohesive sediments of the Lerma River, Mexico. *Revista Internacional de Contaminación Ambiental*, 27(3), 181-190.
- [9] Deepulal, P.M., Kumar, T.R.G., Sujatha, C.H., George, R. (2012). Chemometric study on the trace metal accumulation in the sediments of the Cochin Estuary Southwest coast of India. *Environmental Monitoring and Assessment*, 184,

- 6261–6279. PMID:22083399 [View Article](#) [PubMed/NCBI](#)
- [10]Kong, P., Cheng, X., Sun, R., Chen, L. (2018). The Synergic Characteristics of Surface Water Pollution and Sediment Pollution with Heavy Metals in the Haihe River Basin, Northern China. *Water*, 10,73. [View Article](#)
- [11]Matte, T.D. (2003). Efectos del plomo en la salud de la ni-ez. *Salud Pública Mexicana*, 45(2), 220-224.
- [12]Rainbow, P.S. (1992). The significance of accumulated heavy metal concentrations in marine organisms. In: Miskiewicz AG (Ed). *Assessment of the distribution, impacts and bioaccumulation of contaminants in aquatic environments*, Proceedings of a bioaccumulation Workshop. Sydney. Water Board and Australian Marine Science Association Inc.
- [13]Marín-Guerao, L., Cesar, A., Marín, A., Lloret, J., Vita, R. (2005). Establishing the ecological quality status of soft-bottom mining-impacted coastal water bodies in the scope of Water Framework Directive. *Marine Pollution Bulletin*, 50, 374-387. PMID:15823299 [View Article](#) [PubMed/NCBI](#)
- [14]Acosta, V., C, Lodeiros., W, Senior. (2002). Niveles de metales pesados en sedimentos superficiales en zonas litorales de Venezuela. *Interciencia*, 27, 686-690.
- [15]Rather, B.A., Franson, J.C., Sheffield, S.R., Goddard, C.I., Leonard, N.J., Stand, D., White, P.J. (2008). *Sources and Implications of Lead-based Ammunition and Fishing Tackle on Natural Resources*. The Wildlife Society, Bethesda, MD, USA.
- [16]Zaza, S., de Balogh, K., Palmery, M., et al. (2015). Human exposure in Italy to lead, cadmium and mercury through fish and seafood product consumption from Eastern Central Atlantic Fishing Area. *Journal of Food Composition and Analysis*, 40, 148-153. [View Article](#)
- [17]Signa, G., Mazzola, A., Tramati, C.D., Vizzini, S. (2017). Diet and habitat use influence Hg and Cd transfer to fish consequent biomagnification in highly contaminated area: Augusta Bay (Mediterranean Sea). *Environmental Pollution*, 230, 394-404. PMID:28675849 [View Article](#) [PubMed/NCBI](#)
- [18]Ashraf, A., Saion, E., Gharibshani, E., Yap, C.K. et al. (2018). Distribution of Heavy Metals in Core Marine Sediments of Coastal East Malaysia by Instrumental Neutron Activation Analysis and Inductively Coupled Plasma Spectroscopy. *Applied Radiation and Isotopes*. PMID:29183762 [View Article](#) [PubMed/NCBI](#)
- [19]Liu, Y., Liu, G., Yuan, Z., Liu, H., Lam, P.K.S. (2018). Heavy metals (AS, Hg and V) and stable isotope ratios in fish from Yellow River Estuary, China. *Science of the Total Environment*, 613-614, 462-471. PMID:28918278 [View Article](#) [PubMed/NCBI](#)
- [20]Tong, Y., Wang, M., Bu, X., et al. (2017). Mercury concentrations in China's coastal waters and implications for fish consumption by vulnerable populations. *Environmental pollution*, 231, 396-405. PMID:28818815 [View Article](#) [PubMed/NCBI](#)
- [21]Taylor, D.L., Calabrese, N.M. (2018). Mercury content of blue crabs (*Callinectes Sapidus*) from southern New England coastal habitats: Contamination in an emergent fishery and risks to human consumers. *Marine Pollution Bulletin*, 126, 166-178. PMID:29421084 [View Article](#) [PubMed/NCBI](#)
- [22]Barlas, N., Akbulut, N., Aydogan, M. (2005). Assessment of heavy metal residues in the sediment and water samples of Uluabat Lake, Turkey. *Bulletin of Environmental Contamination and Toxicology*, 74, 286–293. [View Article](#)
- [23]Hall, L.W.J., Scott, M.C., Killen, W.D. (1997). *A Screening Level Probabilistic Ecological Risk Assessment of Copper and Cadmium in the Chesapeake Bay Watershed*. Report. US EPA (Ed), Chesapeake Bay Program office, Annapolis, MD.
- [24]Martínez, G., Rodríguez, L., Senior, W. (2002). Especiación de Cd, Zn, Cr y Pb en núcleos de sedimentos de la Bahía de Bergantín, Estado Anzoátegui, Venezuela. *Interciencia*, 27, 173-178.
- [25]Pedroza, S., Bussy, A., Barceló, I., López, E., Ávila, P. (2010). Organic matter biodegradation and metal behavior in contaminated freshwater sediments. *International Journal of Environmental Technology and Management*, 12(1), 85-94. [View Article](#)
- [26]Zhao, S., Feng, C., Quan, W., et al. (2012). Role of living environments in the accumulation characteristics of heavy metals in fishes and crabs in the Yangtze River Estuary, China. *Marine pollution bulletin*, 64(6), 1163-1171. PMID:22551849 [View Article](#) [PubMed/NCBI](#)
- [27]Sprague, D.D. and Vermaire, J.C. (2018). Legacy Arsenic Pollution of Lakes Near Cobalt, Ontario Canada: Arsenic in Lake Water and Sediment Remains Elevated Nearly a Century After Mining Activity Has Ceased. *Water Air and Soil Pollution*, 229, 87. [View Article](#)
- [28]Bahloul, M., Baati, H., Amdouni, R., Azri, C.

- (2018). Assessment of heavy metals contamination and their potential toxicity in the surface sediments of Sfax solar Saltern, Tunisia. *Environmental Earth Sciences*, 77,27. [View Article](#)
- [29] Urbina-Barreto, I., Hernandez-Valencia, I., Vilamizar, H., Alvarez, M. (2014). Concentraciones de Pb, Cd, Ni, Cr y Cu en Amphimedon Viridis y Niphates erecta (haplosclerida: Nephtyidae) en dos ecosistemas marinos del mar Caribe venezolano. *Sociedad Venezolana de Ecología*, 27(1-2), 31-39.
- [30] Griboff, J., Horacek, M., Wunderlin, D.A., Monferran, M.V. (2018). Bioaccumulation and trophic transfer of metals, As and Se through a freshwater food web affected by atrophic pollution in Córdoba, Argentina. *Ecotoxicology and Environmental Safety*, 148, 275-284. PMID:29078130 [View Article](#) [PubMed/NCBI](#)
- [31] Khan, M. I., Khisroon, M., Khan, A., et al. (2018). Bioaccumulation of Heavy Metals in Water, Sediments, and Tissues and Their Histopathological Effects on Anodonta Cygnea (Linea, 1876) in Kabul river, Khyber Pakhtunkhwa, Pakistan. *Biomed Research International*. [View Article](#)
- [32] Atef El-Taher, Hesham M.H. Zakaly and Reda Elsaman. (2017). Environmental implications and spatial distribution of natural radionuclides and heavy metals in sediments from four harbors in the Egyptian Red Sea coast, *Applied Radiation and Isotopes*.
- [33] Lécivain, N., Aurenche, V., Cottin, N., Frossard, V., Clément, B. (2018). Multi-contamination (heavy metals, polychlorinated biphenyls and polycyclic aromatic hydrocarbons) of littoral sediments and the associated ecological risk assessment in large lake in France (Lake Bourget). *Science of the Total Environment*, 619-620, 854-865. PMID:29734631 [View Article](#) [PubMed/NCBI](#)
- [34] Rajeshkumar, S., Liu, Y., Zhang, X., Ravikumar, B., Bai, G., Li, X. (2017). Studies on seasonal pollution of heavy metals in water, sediment, fish and oyster from the Meiliang Bay of Taihu Lake in China. *Chemosphere*, PMID:29078187
- [35] Moiseenko, T.I. et al. (2018). Ecosystem and human health assessment in relation to aquatic environmental pollution by heavy metals: case study of northwest of the Russian arctic, Kola peninsula. *Environmental Research Letters* in press. [View Article](#)
- [36] Pulich, M. W. (1980). Heavy metals accumulation by selected Halodule Wrightii ASH populations in the Corpus Christi Bay Areas. *Marine Science*, 23, 89-100.
- [37] González-Lozano, M.C., Méndez-Rodríguez, L.C., López-Veneroni, D.G., Vázquez-Botello, A. (2006). Evaluación de la contaminación sedimentos del área portuaria y zona costera de Salina Cruz, Oaxaca, México. *Revista Inter-ciencia*, 31(9), 647-656.
- [38] Celis-Hernandez, O., Rosales-Hoz, L. Cundy, A., Carranza-Edwards. (2017). Sedimentary heavy metal (loid) contamination in the Veracruz shelf, Gulf of Mexico: A baseline survey from a rapidly developing tropical coast. *Marine pollution Bulletin*, 119, 204-213. PMID:28434667 [View Article](#) [PubMed/NCBI](#)
- [39] Botello, A.V., Villanueva, F.S., Rivera, R.F., Velandia, A.L., de la Lanza, G.E. (2018). Analysis and Tendencies of Metals and POPs in Sediment Core from the Alvarado Lagoon System (ALS), Veracruz, Mexico. *Archives of Environmental Contamination and Toxicology*. PMID:29511815 [View Article](#) [PubMed/NCBI](#)
- [40] Aragon, M., Félix, F., García, J., Rivas, D. (2015). Atlas de Sonora de Julio Montané Martí. In Instituto Sonorense de Cultura (Ed).
- [41] Gortares-Moroyoqui, P. (2010). Microbiological Water Quality in Irrigation Water, Treated Wastewater, and Untreated Wastewater and Its Impact on Vegetables in Sonora, Mexico. Doctor of Philosophy Thesis. The University of Arizona. [View Article](#)
- [42] Acosta-Velázquez, J. and A. D. Vázquez-Lule. (2009). Caracterización del sitio de manglar Estero Los Lobos, en Comisión Nacional para el Conocimiento y Uso de la Biodiversidad (CONABIO). Sitios de manglar con relevancia biológica y con necesidades de rehabilitación ecológica. CONABIO, México, D.F.
- [43] Vargas-González, H.H., Arreola-Lizárraga, J.A., García-Hernández, J., et al. (2017). Calidad de sedimentos asociada a actividades antrópicas en lagunas costeras semiáridas subtropicales de la costa central del Golfo de California. *Revista Internacional de Contaminación Ambiental*, 33, 7-22.
- [44] Ortega, P., Vásquez, D. (1992). Distribución y enriquecimiento de metales traen núcleos de sedimento de tres sistemas costeros del estado de Sonora, México. *Acta Oceanográfica del Pacífico*, 7(1), 111-127.
- [45] García-Rico, L., Soto-Cruz, M.S., Jara-Marini, M.E. and Gómez-Álvarez, A. (2004). Fracciones Geoquímicas de Cd, Cu y Pb en sedimentos Costeros Superficiales de Zonas Ostrícolas del Estado de Sonora, México. *Revista Internacional de Contaminación Ambiental*, 20(4), 159-167.

- [46] Norma Oficial Mexicana NOM-059-SEMARNAT-2010. (2010). Protección ambiental -Especies nativas de México de flora y fauna silvestres-Categorías de riesgo y especificaciones para sin inclusión, exclusión o cambio-Lista de especies en riesgo. Diario Oficial de la Federación, 30 de diciembre del 2010.
- [47] Norma Mexicana NMX-AA-051-SCFI-2016. (2016). Análisis de agua-medición de metales-por absorción atómica en aguas naturales, potables, residuales y residuales tratadas. Diario Oficial de la Federación. 7 de diciembre del 2016. In: [View Article](#)
- [48] Norma Mexicana NMX-AA-132-SCFI-2006. (2006). Muestreo de suelos para la identificación y la cuantificación de metales y metaloides, y manejo de la muestra. Secretaría de Economía. 5 de septiembre del 2006. In: [View Article](#)
- [49] Norma Oficial Mexicana NOM-001-ECOL-1996. (1996). Límites máximos permisibles de contaminantes en las descargas de aguas residuales en aguas y bienes nacionales. Secretaría de Medio Ambiente, Recursos Naturales y Pesca. Secretaría de Economía. Diario Oficial de la Federación 6 de enero de 1997. In [View Article](#)
- [50] Plus, M. S. (2001). version 5.1 for Windows. Rockville: Manugistics Group.
- [51] Pérez-Zapata, A.J. (1981). Plomo y mercurio: Lagunas costeras de Tabasco. Centro de Ecodesarrollo (Ed). Unecosistema en peligro (pp. 58-61). México.
- [52] EPA (Environmental Protection Agency). (2013). Aquatic Life Water Quality Criteria Summary. In State of Oregon: Oregon Department of Environmental Quality. [View Article](#)
- [53] CCME (Canadian Council of Ministers of the Environment). (2018). Water and Sediment Quality Guidelines for the Protection of Aquatic Life. In Government of Canada. 20, 71, 124, 131 & chapters = 1, 2, 3, 4, 5 [View Article](#)
- [54] Buchman, M.F. (2008). NOAA Screening Quick Reference Tables, NOAA OR&R Report 08-1. Seattle WA. Office of Response and Restoration Division, National Oceanic and Atmospheric Administration.
- [55] CONAGUA (Comisión Nacional del Agua). (2016). Reporte del clima en México. [View Article](#)